

AZA COMPOUNDS, PHARMACEUTICAL COMPOSITIONS AND
METHODS OF USE

Inventors: Yong-Qian WU
Wei HUANG
Gregory S. HAMILTON

NATH & ASSOCIATES
1030 15th Street NW, 6th Floor
Washington, D.C. 20005
(202) 775-8383

AZA COMPOUNDS, PHARMACEUTICAL COMPOSITIONS AND
METHODS OF USE

5 This application is a Continuation-in-Part of U.S. Patent
Application Serial No. 09/551,618, filed April 17, 2000, which
claims the benefit of U.S. Provisional Application No.
60/164,950 filed November 12, 1999.

BACKGROUND OF THE INVENTION

Field of Invention

10 The present invention relates generally to N-substituted
cyclic aza compounds, pharmaceutical compositions comprising
such compounds, and their preparation and use for preventing
and/or treating neurological disorders; for treating alopecia
and promoting hair growth; for treating vision disorders
15 and/or improving vision; for treating memory impairment and/or
enhancing memory performance; and for preventing or treating
hearing loss in an animal.

Description of Related Art

20 It has been found that picomolar concentrations of an
immunosuppressant such as FK506 and rapamycin stimulate
neurite outgrowth in PC12 cells and sensory nervous, namely
dorsal root ganglion cells (DRGs). Lyons et al., *Proc. Of*
Natl. Acad. Sci., 1994 vol. 91, pp. 3191-3195. In whole
animal experiments, FK506 has been shown to stimulate nerve
25 regeneration following facial nerve injury and results in
functional recovery in animals with sciatic nerve lesions.

30 Several neurotrophic factors effecting specific neuronal
populations in the central nervous system have been
identified. For example, it has been hypothesized that
Alzheimer's disease results from a decrease or loss of nerve
growth factor (NGF). It has thus been proposed to treat
Alzheimer's patients with exogenous nerve growth factor or
other neurotrophic proteins such as brain derived nerve factor

(BDNF), glial derived nerve factor, ciliary neurotrophic factor, and neurotrophin-3 to increase the survival of degenerating neuronal populations.

Clinical application of these proteins in various neurological disease states is hampered by difficulties in the delivery and bioavailability of large proteins to nervous system targets. By contrast, immunosuppressant drugs with neurotrophic activity are relatively small and display excellent bioavailability and specificity. However, when administered chronically, immunosuppressants exhibit a number of potentially serious side effects including nephrotoxicity, such as impairment of glomerular filtration and irreversible interstitial fibrosis (Kopp et al., 1991, *J. Am. Soc. Nephrol.* 1:162); neurological deficits, such as involuntary tremors, or non-specific cerebral angina such as non-localized headaches (De Groen et al., 1987, *N. Engl. J. Med.* 317:861); and vascular hypertension with complications resulting therefrom (Kahan et al., 1989 *N. Engl. J. Med.* 321: 1725).

Accordingly, there is a need for non-immunosuppressive, small-molecule compounds which are useful for neurotrophic effects and for treating neurodegenerative disorders.

Hair loss occurs in a variety of situations. These situations include male pattern alopecia, alopecia senilis, alopecia areata, diseases accompanied by basic skin lesions or tumors, and systematic disorders such as nutritional disorders and internal secretion disorders. The mechanisms causing hair loss are very complicated, but in some instances can be attributed to aging, genetic disposition, the activation of male hormones, the loss of blood supply to hair follicles, and scalp abnormalities.

The immunosuppressant drugs FK506, rapamycin and cyclosporin are well known as potent T-cell specific

immunosuppressants, and are effective against graft rejection after organ transplantation. It has been reported that topical, but not oral, application of FK506 (Yamamoto et al., *J. Invest. Dermatol.*, 1994, 102, 160-164; Jiang et al., *J. Invest. Dermatol.* 1995, 104, 523-525) and cyclosporin (Iwabuchi et al., *J. Dermatol. Sci.* 1995, 9, 64-69) stimulates hair growth in a dose-dependent manner. One form of hair loss, alopecia areata, is known to be associated with autoimmune activities; hence, topically administered immunomodulatory compounds are expected to demonstrate efficacy for treating that type of hair loss. The hair growth stimulating effects of FK506 have been the subject of an international patent filing covering FK506 and structures related thereto for hair growth stimulation (Honbo et al., EP 0 423 714 A2). Honbo et al. discloses the use of relatively large tricyclic compounds, known for their immunosuppressive effects, as hair revitalizing agents.

The hair growth and revitalization effects of FK506 and related agents are disclosed in many U.S. patents (Goulet et al., U.S. Patent No. 5,258,389; Luly et al., U.S. Patent No. 5,457,111; Goulet et al., U.S. Patent No. 5,532,248; Goulet et al., U.S. Patent No. 5,189,042; and Ok et al., U.S. Patent No. 5,208,241; Rupprecht et al., U.S. Patent No. 5,284,840; Organ et al., U.S. Patent No. 5,284,877). These patents claim FK506 related compounds. Although they do not claim methods of hair revitalization, they disclose the known use of FK506 for effecting hair growth. Similar to FK506 (and the claimed variations in the Honbo et al. patent), the compounds claimed in these patents are relatively large. Further, the cited patents relate to immunomodulatory compounds for use in autoimmune related diseases, for which FK506's efficacy is well known.

Other U.S. patents disclose the use of cyclosporin and related compounds for hair revitalization (Hauer et al., U.S. Patent No. 5,342,625; Eberle, U.S. Patent No. 5,284,826; Hewitt et al., U.S. Patent No. 4,996,193). These patents also
5 relate to compounds useful for treating autoimmune diseases and cite the known use of cyclosporin and related immunosuppressive compounds for hair growth.

However, immunosuppressive compounds by definition suppress the immune system and also exhibit other toxic side
10 effects. Accordingly, there is a need for non-immunosuppressive, small molecule compounds which are useful as hair revitalizing compounds.

The visual system is composed of the eyes, ocular adnexa and the visual pathways. Dysfunction of the visual system may
15 lead to permanent or temporary visual impairment, i.e. a deviation from normal in one or more functions of the eye. Visual impairment manifests itself in various ways and includes a broad range of visual dysfunctions and disturbances. Without limitation, these dysfunctions and
20 disturbances include partial or total loss of vision, the need for correction of visual acuity for objects near and far, loss of visual field, impaired ocular motility without diplopia (double vision), impaired or skewed color perception, limited adaptation to light and dark, diminished accommodation, metamorphopsic distortion, impaired binocular vision, paresis
25 of accommodation, iridoplegia, entropion, ectropion, epiphora, lagophthalmos, and scarring. See *Physicians' Desk Reference (PDR) for Ophthalmology*, 16th Edition, 6:47 (1988). The visual system may be adversely affected by various
30 ophthalmologic disorders, diseases, injuries, and complications, including, without limitation, genetic disorders; disorders associated with aging or degenerative

diseases; disorders correlating to physical injury to the eye, head, or other parts of the body resulting from external forces; disorders resulting from environmental factors; disorders resulting from a broad range of diseases; and combinations of any of the above.

The visual system is a complex system composed of numerous components. Visual impairment can involve the entire visual system, any one component, or any combination of components, depending upon the precise nature of the circumstances. The eye is composed of a lens, which is suspended in the zonules of Zinn and is focused by the ciliary body. The ciliary body also secretes aqueous humor, which fills the posterior chamber, passes through the pupil into the anterior chamber, then drains primarily via the canal of Schlemm. The iris regulates the quantity of light entering the eye by adjusting the size of its central opening, the pupil. A visual image is focused onto the retina, the fovea centralis being the retinal area of sharpest visual acuity. The conjunctiva is the mucus membrane which lines the eyelids and the eyeball, and ends abruptly at the limbus conjunctivae, the edge of the conjunctiva overlapping the cornea. The cornea is the clear, transparent anterior portion of the fibrous coat of the eye; it is important in light refraction and is covered with an epithelium that differs in many respects from the conjunctival epithelium.

The retina is the innermost, light sensitive portion of the eye, containing two types of photoreceptors, cones, which are responsible for color vision in brighter light, and rods, which are essential for vision in dim light but do not perceive colors. After light passes through the cornea, lens system, and the vitreous humor, it enters the retina from the inside; that is, it passes through the ganglion cells and

nerve fibers, the inner and outer plexiform layers, the inner and outer nuclear layers, and the internal and external limiting membranes before it finally reaches the layer of photoreceptors located near the outside of the retina, just inside the outermost pigment epithelium layer. The cells of the pigment epithelium layer act as an anatomical barrier to liquids and substances located outside of the eye, forming the "blood-retina" barrier, and provide nourishment, oxygen, a source of functionally useful substances like vitamin A, and phagocytosis of decomposition products to photoreceptor cells. There is no anatomical connection between the pigment epithelium and the photoreceptor layer, permitting separation of the layers in some pathological situations.

When rods or cones are excited by light, signals are transmitted through successive neurons in the retina itself, into the optic nerve fibers, and ultimately to the cerebral cortex. Both rods and cones contain molecules that decompose on exposure to light and, in the process, excite the nerve fibers leading from the eye. The molecule in rods is rhodopsin. The three light-sensitive molecules in cones, collectively called iodopsin, have compositions only slightly different from that of rhodopsin and are maximally excited by red, blue, or green light, respectively.

Neither rods nor cones generate action potentials. Rather, the light-induced membrane hyperpolarization generated in the outer, photosensitive segment of a rod or cone cell is transmitted from the outer segment through the inner segment to the synaptic body by direct conduction of the electrical voltage itself, a process called electrotonic conduction. At the synaptic body, the membrane potential controls the release of an unknown transmitter molecule. In low light, rod and cone cell membranes are depolarized and the rate of

transmitter release is greatest. Light-induced hyperpolarization causes a marked decrease in the release of transmitter molecules.

5 The transmitters released by rod and cone cells induce signals in the bipolar neurons and horizontal cells. The signals in both these cells are also transmitted by electrotonic conduction and not by action potential.

10 The rod bipolar neurons connect with as many as 50 rod cells, while the dwarf and diffuse bipolar cells connect with one or several cone cells. A depolarizing bipolar cell is stimulated when its connecting rods or cones are exposed to light. The release of transmitter molecules inhibits the depolarizing bipolar cell. Therefore, in the dark, when the rods and cones are secreting large quantities of transmitter
15 molecules, the depolarizing bipolar cells are inhibited. In the light, the decrease in release of transmitter molecules from the rods and cones reduces the inhibition of the bipolar cell, allowing it to become excited. In this manner, both positive and negative signals can be transmitted through
20 different bipolar cells from the rods and cones to the amacrine and ganglion cells.

As their name suggests, horizontal cells project horizontally in the retina, where they may synapse with rods, cones, other horizontal cells, or a combination of cells
25 types. The function of horizontal cells is unclear, although some mechanism in the convergence of photoreceptor signaling has been postulated.

All types of bipolar cells connect with ganglion cells, which are of two primary types. A-type ganglion cells
30 predominately connect with rod bipolar cells, while B-type ganglion cells predominately connect with dwarf and diffuse bipolar cells. It appears that A-type ganglion cells are

sensitive to contrast, light intensity, and perception of movement, while B-type ganglion cells appear more concerned with color vision and visual acuity.

5 Like horizontal cells, the Amacrine cells horizontally synapse with several to many other cells, in this case bipolar cells, ganglion cells, and other Amacrine cells. The function of Amacrine cells is also unclear.

10 The axons of ganglion cells carry signals into the nerve fiber layer of the eye, where the axons converge into fibers which further converge at the optic disc, where they exit the eye as the optic nerve. The ganglion cells transmit their signals through the optic nerve fibers to the brain in the form of action potentials. These cells, even when
15 unstimulated, transmit continuous nerve impulses at an average, baseline rate of about 5 per second. The visual signal is superimposed onto this baseline level of ganglion cell stimulation. It can be either an excitatory signal, with the number of impulses increasing above the baseline rate, or an inhibitory signal, with the number of nerve impulses
20 decreasing below the baseline rate.

As part of the central nervous system, the eye is in some ways an extension of the brain; as such, it has a limited capacity for regeneration. This limited regeneration capacity further complicates the challenging task of improving vision,
25 resolving dysfunction of the visual system, and/or treating or preventing ophthalmologic disorders. Many disorders of the eye, such as retinal photic injury, retinal ischemia-induced eye injury, age-related macular degeneration, free radical-induced eye diseases, as well as numerous other disorders, are
30 considered to be entirely untreatable. Other ophthalmologic disorders, e.g., disorders causing permanent visual impairment, are corrected only by the use of ophthalmic

devices and/or surgery, with varying degrees of success.

The immunosuppressant drugs FK506, rapamycin, and cyclosporin are well known as potent T-cell specific immunosuppressants, and are effective against autoimmunity, transplant or graft rejection, inflammation, allergic responses, other autoimmune or immune-mediated diseases, and infectious diseases. It has been disclosed that application of Cyclosporin, FK-506, Rapamycin, Buspirone, Spiperone, and/or their derivatives are effective in treating some ophthalmologic disorders of these types. Several ophthalmologic disorders or vision problems are known to be associated with autoimmune and immunologically-mediated activities; hence, immunomodulatory compounds are expected to demonstrate efficacy for treating those types of ophthalmologic disorders or vision problems.

The effects of FK506, Rapamycin, and related agents in the treatment of ophthalmologic diseases are disclosed in several U.S. patents (Goulet et al., U.S. Patent No. 5,532,248; Mochizuki et al., U.S. Patent No. 5,514,686; Luly et al., U.S. Patent No. 5,457,111; Russo et al., U.S. Patent No. 5,441,937; Kulkarni, U.S. Patent No. 5,387,589; Asakura et al., U.S. Patent No. 5,368,865; Goulet et al., U.S. Patent No. 5,258,389; Armistead et al., U.S. Patent No. 5,192,773; Goulet et al., U.S. Patent No. 5,189,042; and Fehr, U.S. Patent No. 5,011,844). These patents claim FK506 or Rapamycin related compounds and disclose the known use of FK506 or Rapamycin related compounds in the treatment of ophthalmologic disorders in association with the known immunosuppressive effects of FK506 and Rapamycin. The compounds disclosed in these patents are relatively large. Further, the cited patents relate to immunomodulatory compounds limited to treating autoimmunity or related diseases, or immunologically-

mediated diseases, for which the efficacy of FK506 and Rapamycin is well known.

Other U.S. patents disclose the use of cyclosporin, Spiperone, Buspirone, their derivatives, and other immunosuppressive compounds for use in the treatment of ophthalmologic diseases (Sharpe et al., U.S. Patent No. 5,703,088; Sharpe et al., U.S. Patent No. 5,693,645; Sullivan, U.S. Patent No. 5,688,765; Sullivan, U.S. Patent No. 5,620,921; Sharpe et al., U.S. Patent No. 5,574,041; Eberle, U.S. Patent No. 5,284,826; Sharpe et al., U.S. Patent No. 5,244,902; Chiou et al., U.S. Patent Nos. 5,198,454 and 5,194,434; and Kaswan, U.S. Patent No. 4,839,342). These patents also relate to compounds useful for treating autoimmune diseases and cite the known use of cyclosporin, Spiperone, Buspirone, their derivatives, and other immunosuppressive compounds in treating ocular inflammation and other immunologically-mediated ophthalmologic diseases.

The immunosuppressive compounds disclosed in the prior art suppress the immune system, by definition, and also exhibit other toxic side effects. Accordingly, there is a need for non-immunosuppressant, small molecule compounds, and compositions and methods for use of such compounds, that are useful in improving vision; preventing, treating, and/or repairing visual impairment or dysfunction of the visual system; and preventing, treating, and/or resolving ophthalmologic disorders.

There are also a number of patents on non-immunosuppressive compounds disclosing methods of use for permitting or promoting wound healing (whether from injury or surgery); controlling intraocular pressure (often resulting from glaucoma); controlling neurodegenerative eye disorders, including damage or injury to retinal neurons, damage or

injury to retinal ganglion cells, and macular degeneration; stimulating neurite outgrowth; preventing or reducing oxidative damage caused by free radicals; and treating impaired oxygen and nutrient supply, as well as impaired waste product removal, resulting from low blood flow. These non-immunosuppressive substances fall into one of two general categories: naturally occurring molecules, such as proteins, glycoproteins, peptides, hormones, and growth factors; and synthetic molecules.

Within the group of naturally occurring non-immunosuppressive molecules, several hormones, growth factors, and signaling molecules have been patented for use as supplements to naturally occurring quantities of such molecules, as well as for targeting of specific cells where the particular molecule does not naturally occur in a mature individual. These patents generally claim methods of use for reducing or preventing the symptoms of ocular disease, or arresting or reversing vision loss.

Specifically, Louis et al., U.S. Patent Nos. 5,736,516 and 5,641,749, disclose the use of a glial cell line derived neurotrophic factor (GDNF) to stop or reverse the degeneration of retinal neurons (i.e. photoreceptors) and retinal ganglion cells caused by glaucoma, or other degenerative or traumatic retinal diseases or injuries. O'Brien, et al., U.S. Patent Nos. 5,714,459 and 5,700,909, disclose the use of a glycoprotein, Saposin, and its derivatives for stimulating neurite outgrowth and increasing myelination. To stop or reverse degeneration of retinal neurons, LaVail et al., U.S. Patent No. 5,667,968, discloses the use of a variety of neurotrophic proteins, including brain-derived neurotrophic factor, ciliary neurotrophic factor, neurotrophin-3 or neurotrophin-4, acidic or basic fibroblast growth factors,

interleukin, tumor necrosis factor- α , insulin-like growth factor-2 and other growth factors. Wong et al., U.S. Patent No. 5,632,984, discloses the use of interferons, especially interferon α -2a, for treating the symptoms of macular degeneration by reducing hemorrhage and limiting neovascularization. Finally, Wallace et al., U.S. Patent No. 5,441,937, discloses the use of a lung-derived neurotrophic factor (NTF) to maintain the functionality of ciliary ganglion and parasympathetic neuron cells.

A key characteristic of factors derived from specific cell lines is their localization to specific cell lines or tissues; systemic treatment with these molecules would run a substantial risk of unintended, and potentially dangerous, effects in cell lines where the genes encoding these molecules are inactive. Similarly, hormones and growth factors often activate a large number of genes in many cell lines; again, non-localized application of these molecules would run a substantial risk of provoking an inappropriate, and potentially dangerous, response.

Within the category of synthetic molecules, most of the patented compounds are immunosuppressive and disclose uses in treating inflammatory, autoimmune, and allergic responses, as discussed above. A few others are non-immunosuppressive and claim the ability to treat cellular degeneration, and in some cases promote cellular regeneration, most often in the context of their antioxidant properties.

Specifically, Tso et al., U.S. Patent No. 5,527,533, discloses the use of astaxanthin, a carotenoid antioxidant, for preventing or reducing photoreceptor damage resulting from the presence of free radicals. Similarly, Babcock et al., U.S. Patent No. 5,252,319, discloses the use of antioxidant aminosteroids for treating eye disease and injury, by

increasing resistance to oxidative damage. Freeman, U.S. Patent No. 5,468,752, discloses the use of the antiviral phosphonylmethoxyalkylcytosines to reduce abnormally increased intraocular pressure.

5 Naturally occurring hormones, growth factors, cytokines, and signaling molecules are generally multifunctional and activate many genes in diverse cell lines. The present compounds do not, thus avoiding the unexpected, and potentially dangerous, side effects of systemic use.
10 Similarly, the present compounds also avoid the potential unexpected side effects of introducing cell line-specific molecules into other cell lines where they do not naturally occur.

The epithelial hair cells in the organ of Corti of the
15 inner ear, transduce sound into neural activity, which is transmitted along the cochlear division of the eighth cranial nerve. This nerve consists of fibers from three types of neurons (Spoendllin, H. H., in Friedmann, I. Ballantyne, J., eds. "Ultrastructural Atlas of the Inner Ear", London, Butterworth, pp. 133-164, (1984)) 1) afferent neurons, which
20 lie in the spiral ganglion and connect the cochlea to the brainstem; 2) efferent olivocochlear neurons, which originate in the superior olivary complex; and 3) autonomic adrenergic neurons, which originate in the cervical sympathetic trunk and innervate the cochlea. In the human, there are approximately
25 30,000 afferent cochlear neurons, with myelinated axons, each consisting of about 50 lamellae, and 4-6 μ m in diameter. This histologic structure forms the basis of uniform conduction velocity, which is an important functional feature.
30 Throughout the length of the auditory nerve, there is a trophic arrangement of afferent fibers, with 'basal' fibers wrapped over the centrally placed 'apical' fibers in a twisted

rope-like fashion. Spoendlin (Spoendlin, H. H. in Naunton, R. F., Fernadex, C. eds., "Evoked Electrical Activity in the Auditory Nervous System", London, Academic Press, pp. 21-39, (1978)) identified two types of afferent neurons in the spiral ganglion on the basis of morphologic differences: type I cells (95%) are bipolar and have myelinated cell bodies and axons that project to the inner hair cells. Type II cells (5%) are monopolar with unmyelinated axons and project to the outer hair cells of the organ of Corti. Each inner hair cell is innervated by about 20 fibers, each of which synapses on only one cell. In contrast, each outer hair cell is innervated by approximately six fibers, and each fiber branches to supply approximately 10 cells. Within the cochlea, the fibers divide into: 1) an inner spiral group, which arises primarily ipsilaterally and synapses with the afferent neurons to the inner hair cells, and 2) a more numerous outer radial group, which arises mainly contralaterally and synapses directly with outer hair cells. There is a minimal threshold at one frequency, the characteristic or best frequency, but the threshold rises sharply for frequencies above and below this level (Pickles, J.O. in "Introduction to the Physiology of Hearing", London, Academic Press, pp. 71-106, (1982)). Single auditory nerve fibers therefore appear to behave as band-pass filters. The basilar membrane vibrates preferentially to different frequencies, at different distances along its length, and the frequency selectivity of each cochlear nerve fiber is similar to that of the inner hair cell to which the fiber is connected. Thus, each cochlear nerve fiber exhibits a tuning curve covering a different range of frequencies from its neighboring fiber (Evans, E.F. in Beagley H.A. ed., "Auditory investigation: The Scientific and Technological basis", New York, Oxford University Pressm (1979)). By this

mechanism, complex sounds are broken down into component frequencies (frequency resolution) by the filters of the inner ear.

5 Impairment anywhere along the auditory pathway, from the
external auditory canal to the central nervous system, may
result in hearing loss. The auditory apparatus can be
subdivided into the external and middle ear, inner ear and
auditory nerve and central auditory pathways. Auditory
10 information in humans is transduced from a mechanical signal
to a neurally conducted electrical impulse by the action of
approximately 15,000 epithelial cells (hair cells) and 30,000
first-order neurons (spiral ganglion cells) in the inner ear.
All central fibers of spiral ganglion neurons form synapses in
the cochlear nucleus of the pontine brainstem, The number of
15 neurons involved in hearing increases dramatically from the
cochlea to the auditory brain stem and the auditory cortex.
All auditory information is transduced by only 15,000 hair
cells, of which the so-called inner hair cells, numbering
3500, are critically important, since they form synapses with
20 approximately 90 percent of the 30,000 primary auditory
neurons. Thus, damage to a relatively few cells in the
auditory periphery can lead to substantial hearing loss.
Hence, most causes of sensorineural loss can be ascribed to
lesions in the inner ear (Nadol, J.B., New England Journal of
25 Medicine, (1993), 329:1092-1102).

 Hearing loss can be on the level of conductivity,
sensorineural and central level. Conductive hearing loss is
caused by lesions involving the external or middle ear,
resulting in the destruction of the normal pathway of airborne
30 sound amplified by the tympanic membrane and the ossicles to
the inner ear fluids. Sensorineural hearing loss is due to
lesions of the central auditory pathways. These consist of

the cochlear and dorsal olivary nucleus complex, inferior colliculi, medial geniculate bodies, auditory cortex in the temporal lobes and interconnecting afferent and efferent fiber tracts (Adams R.D. and Maurice, V., eds., in "Principles of Neurology", (1989), McGraw-Hill Information Services Company, pp. 226-246).

Trauma due to acoustic overstimulation is another leading cause of deafness. There is individual susceptibility to trauma from noise. Clinically important sensorineural hearing loss may occur in some people exposed to high-intensity noise, even below levels approved by the Occupational Safety and Health Agency (Osguthorpe, J.D., ed., Washington D.C., American Academy of Otolaryngology-Head and Neck Surgery Foundation, (1988)).

Demyelinating processes, such as multiple sclerosis, may cause sensorineural hearing loss (Noffsinger, D., et al., Acto Otolaryngol. Suppl. (Stockh.) (1972), 303:1-63). More recently, a form of immune-mediated sensorineural hearing loss has been recognized (McCabe, B.F., Ann. Otol. Rhinol. Laryngol. (1979), 88:585-9). The hearing loss is usually bilateral, is rapidly progressive (measured in weeks and months), and may or may not be associated with vestibular symptoms.

A variety of tumors, both primary and metastatic, can produce either a conductive hearing loss, or a sensorineural hearing loss, by invading the inner ear or auditory nerve (Houck, J.R., et al., Otolaryngol. Head Neck Surg. (1992), 106:92-7). A variety of degenerative disorders of unknown cause can produce sensorineural hearing loss. Meniere's syndrome (Nadol, J.B., ed., "Meniere's Disease: Pathogenesis, Pathophysiology, Diagnosis, And Treatment," Amsterdam: Kugler & Ghedini (1989)), characterized by fluctuating sensorineural

hearing loss, episodic vertigo, and tinnitus, appears to be caused by a disorder of fluid homeostasis within the inner ear, although the pathogenesis remains unknown. Sudden idiopathic sensorineural hearing loss (Wilson, W.R., et al., Arch. Otolaryngol. (1980), 106:772-6), causing moderate-to-severe sensorineural deafness, may be due to various causes, including inner ear ischemia and viral labyrinthitis.

Regardless of the cause, there exists a need to prevent or treat sensorineural hearing loss. The present invention provides such a method.

SUMMARY OF THE INVENTION

The present invention is directed to novel N-substituted cyclic aza compounds, and their preparation and use for treating neurodegenerative disorders, for treating alopecia and related hair loss disorders, for treating vision disorders and/or improving vision, for treating memory impairment and/or enhancing memory performance, and for treating sensorineural hearing loss. Preferred compounds include N-diketo cyclic aza derivative compounds, N-sulfonyl cyclic aza derivative compounds, tertiary N-aminocarbonyl cyclic aza compounds, and secondary N-aminocarbonyl cyclic aza compounds, wherein the diketo, sulfonyl, or aminocarbonyl group is attached to one of the nitrogen atoms (the 1-nitrogen) of the central aza heterocyclic ring, with an additional substituent substituted onto the 2-nitrogen of the central aza heterocyclic ring. These novel classes of aza derivative compounds contain the additional feature of a 5-7 membered aza heterocyclic ring in the central structure.

These compounds stimulate neuronal regeneration and outgrowth and as such are useful for treating neurological disorders and neurodegenerative diseases. These compounds also promote hair growth and as such are useful for treating


5

10



n is 1-3;

15



wherein said R_1 group is either unsubstituted or additionally substituted with R_3 ;


R_2 is selected from the group consisting of hydrogen, C_1 - C_9 , straight or branched chain alkyl, C_2 - C_9 , straight or branched chain alkenyl, C_2 - C_9 , straight or branched chain alkynyl, aryl,

5 R₃ is selected from the group consisting of hydrogen, C₁-
C₉ alkyl, C₂-C₉ straight or branched chain alkenyl, C₂-C₉
straight or branched chain alkynyl, C₁-C₉ alkoxy, C₂-C₉
alkenyloxy, aryloxy, phenoxy, benzyloxy, hydroxy, carboxy, C₁-
C₉ thioalkyl, C₂-C₉ thioalkenyl, C₁-C₉ alkylamino, C₂-C₉
10 alkenylamino, cyano, nitro, imino, sulfonyl, thiocarbonyl,
sulfhydryl, halo, haloalkyl, trifluoromethyl, aryl,
heteroaryl, carbocycle, and heterocycle,

X is 0 or S.


25 or a pharmaceutically acceptable salt, ester or solvate thereof, wherein:

R₁ is selected from the group consisting of -CR₃, -COOR₃, -COR₃, -COOH, -SO₃H, -SO₂HNR₃, -PO₂(R₃)₂, -CN, -PO₃(R₃)₂, -OR₃, -



II

SR_3 , $-NHCOR_3$, $-N(R_3)_2$, $-CON(R_3)_2$, $-CONH(O)R_3$, $-CONHNHSO_2R_3$, $-COHNSO_2R_3$, $-CONR_3CN$,




- 5 wherein said R_1 group is either unsubstituted or additionally substituted with R_3 ;

R_2 is selected from the group consisting of hydrogen, C_1 - C_9 straight or branched chain alkyl, C_2 - C_9 straight or branched chain alkenyl, C_2 - C_9 straight or branched chain alkynyl, aryl, heteroaryl, carbocycle, or heterocycle, wherein said alkyl, alkenyl, alkynyl, aryl, heteroaryl, carbocycle, or heterocycle is unsubstituted or substituted with one or more substituents selected from R_3 ; and

R_3 is selected from the group consisting of hydrogen, C_1 - C_9 alkyl, C_2 - C_9 straight or branched chain alkenyl, C_2 - C_9 straight or branched chain alkynyl, C_1 - C_9 alkoxy, C_2 - C_9 alkenyloxy, aryloxy, phenoxy, benzyloxy, hydroxy, carboxy, C_1 - C_9 thioalkyl, C_2 - C_9 thioalkenyl, C_1 - C_9 alkylamino, C_2 - C_9 alkenylamino, cyano, nitro, imino, sulfonyl, thiocarbonyl, sulfhydryl, halo, haloalkyl, trifluoromethyl, aryl, heteroaryl, carbocycle, and heterocycle,

wherein said alkyl, alkenyl, alkynyl, alkoxy, alkenyloxy, aryloxy, thioalkyl, thioalkenyl, alkylamino, alkenylamino, aryl, heteroaryl, carbocycle, or heterocycle group is optionally substituted with a hydroxy, carboxy, carbonyl, cyano, nitro, imino, sulfonyl, thiocarbonyl, sulfhydryl, halo, haloalkyl, trifluoromethyl, aryl, heteroaryl, carbocycle, or heterocycle group.

Yet another preferred embodiment of this invention is a compound of formula III:




III

or a pharmaceutically acceptable salt, ester or solvate thereof, wherein:

n is 1-3;

R_1 is selected from the group consisting of $-CR_3$, $-COOR_3$,
 $-COR_3$, $-COOH$, $-SO_3H$, $-SO_2HNR_3$, $-PO_2(R_3)_2$, $-CN$, $-PO_3(R_3)_2$, $-OR_3$, $-$
 SR_3 , $-NHCOR_3$, $-N(R_3)_2$, $-CON(R_3)_2$, $-CONH(O)R_3$, $-CONHNHSO_2R_3$, $-$
 5 $COHNSO_2R_3$, $-CONR_3CN$,



wherein said R_1 group is either unsubstituted or additionally substituted with R_3 ;


5 R and R_2 are independently C_1 - C_9 alkyl, C_2 - C_9 alkenyl, aryl, heteroaryl, carbocycle, or heterocycle, wherein said alkyl, alkenyl, aryl, heteroaryl, carbocycle, or heterocycle

is unsubstituted or substituted with one or more substituent(s) selected from R_3 ; and

R_3 is selected from the group consisting of hydrogen, C_1 - C_9 alkyl, C_2 - C_9 straight or branched chain alkenyl, C_2 - C_9 straight or branched chain alkynyl, C_1 - C_9 alkoxy, C_2 - C_9 alkenyloxy, aryloxy, phenoxy, benzyloxy, hydroxy, carboxy, C_1 - C_9 thioalkyl, C_2 - C_9 thioalkenyl, C_1 - C_9 alkylamino, C_2 - C_9 alkenylamino, cyano, nitro, imino, sulfonyl, thiocarbonyl, sulfhydryl, halo, haloalkyl, trifluoromethyl, aryl, heteroaryl, carbocycle, and heterocycle,

wherein said alkyl, alkenyl, alkynyl, alkoxy, alkenyloxy, aryloxy, thioalkyl, thioalkenyl, alkylamino, alkenylamino, aryl, heteroaryl, carbocycle, or heterocycle group is optionally substituted with a hydroxy, carboxy, carbonyl, cyano, nitro, imino, sulfonyl, thiocarbonyl, sulfhydryl, halo, haloalkyl, trifluoromethyl, aryl, heteroaryl, carbocycle, or heterocycle group.


Yet another preferred embodiment of this invention is a compound of formula IV:



or a pharmaceutically acceptable salt, ester or solvate thereof, wherein:

n is 1-3;

R_1 is selected from the group consisting of $-CR_3$, $-COOR_3$, $-COR_3$, $-COOH$, $-SO_3H$, $-SO_2HNR_3$, $-PO_2(R_3)_2$, $-CN$, $-PO_3(R_3)_2$, $-OR_3$, SR_3 , $-NHCOR_3$, $-N(R_3)_2$, $-CON(R_3)_2$, $-CONH(O)R_3$, $-CONHNHSO_2R_3$, $-COHNSO_2R_3$, $-CONR_3CN$,



wherein said R₁ group is either unsubstituted or additionally substituted with R₃; and

5 R₂ is C₁-C₉ alkyl, C₂-C₉ alkenyl, aryl, heteroaryl, carbocycle, or heterocycle, wherein said alkyl, alkenyl, aryl,

heteroaryl, carbocycle, or heterocycle is substituted with one or more substituent(s) selected from R₃; and

R₃ is selected from the group consisting of hydrogen, C₁-C₉ alkyl, C₂-C₉ straight or branched chain alkenyl, C₂-C₉ straight or branched chain alkynyl, C₁-C₉ alkoxy, C₂-C₉ alkenyloxy, aryloxy, phenoxy, benzyloxy, hydroxy, carboxy, C₁-C₉ thioalkyl, C₂-C₉ thioalkenyl, C₁-C₉ alkylamino, C₂-C₉ alkenylamino, cyano, nitro, imino, sulfonyl, thiocarbonyl, sulfhydryl, halo, haloalkyl, trifluoromethyl, aryl, heteroaryl, carbocycle, and heterocycle,

wherein said alkyl, alkenyl, alkynyl, alkoxy, alkenyloxy, aryloxy, thioalkyl, thioalkenyl, alkylamino, alkenylamino, aryl, heteroaryl, carbocycle, or heterocycle group is optionally substituted with a hydroxy, carboxy, carbonyl, cyano, nitro, imino, sulfonyl, thiocarbonyl, sulfhydryl, halo, haloalkyl, trifluoromethyl, aryl, heteroaryl, carbocycle, or heterocycle group.

Another preferred embodiment is a pharmaceutical composition, comprising:

- (i) an effective amount of the compound of formulae I, II, III, or IV; and
- (ii) a pharmaceutically acceptable carrier.

For neurotrophic compositions, a neurotrophic factor different from the present inventive compounds may also be administered or otherwise included in the composition.

Another preferred embodiment of the invention is a method of treating a neurological disorder in an animal, comprising administering to the animal an effective amount of the compound of formulae I, II, III, or IV to stimulate growth of damaged peripheral nerves or to promote neuronal regeneration.

Another preferred embodiment of the invention is a method of stimulating growth of damaged peripheral nerves, comprising administering to a damaged peripheral nerve an effective amount of the compound of formulae I, II, III, or IV to stimulate or promote growth of the damaged peripheral nerve.

Another preferred embodiment of the invention is a method of promoting neuronal regeneration and growth in animals, comprising administering to an animal an effective amount of the compound of formulae I, II, III, or IV to promote neuronal regeneration.

Another preferred embodiment of the invention is a method of preventing neurodegeneration in animals, comprising administering to an animal an effective amount of the compound of formulae I, II, III, or IV to prevent neurodegeneration.

Another preferred embodiment of the invention is a method for treating alopecia or promoting hair growth in an animal, comprising administering to an animal an effective amount of the compound of formulae I, II, III, or IV.

Another preferred embodiment of the invention is a method for treating a vision disorder, improving vision, treating memory impairment, enhancing memory performance, or treating sensorineural hearing loss in an animal, comprising administering to an animal an effective amount of the compound of formulae I, II, III, or IV.

The present invention further contemplates a process for preparing the cyclic aza derivative compounds of the invention.

The present invention further contemplates the compound(s) of the invention for use in treatment of a disease. In particular, the present invention contemplates the compound(s) of the invention for use in treatment of the disorders enumerated herein.

The present invention further contemplates the compound(s) of the invention for use in the preparation of a medicament or pharmaceutical composition. In particular, the invention contemplates the compound(s) of the invention for use in the preparation of a medicament or pharmaceutical composition for treatment of the disorders enumerated herein.

The invention also provides for the use of compound(s) of the invention for treating a disease. In particular, the invention provides for the use of compound(s) of the invention for treating the disorders enumerated herein.

The invention also provides for the use of compound(s) of the invention in the manufacture of a medicament or pharmaceutical composition. In particular, the invention provides for the use of compound(s) of the invention in the manufacture of a medicament or pharmaceutical composition for the treatment of the disorders enumerated herein. Such pharmaceutical compositions include, as appropriate to the specific disorder, topical, systemic, oral, or injectable formulations. It is further contemplated that the compound(s) of the invention may be administered with an effective amount of a second therapeutic agent for the treatment of the enumerated disorders. A variety of pharmaceutical formulations and different delivery techniques are described in further detail below.

DETAILED DESCRIPTION OF THE INVENTION

Definitions

"Alkyl", alone or in combination, refers to a straight-chain or branched-chain hydrocarbon radical containing a designated number of carbon atoms. For example, C₁-C₉ alkyl is a straight or branched hydrocarbon chain containing 1 to 9 carbon atoms. Examples of such radicals include methyl, ethyl, propyl, iso-propyl, butyl, iso-butyl, tert-butyl, n-pentyl, n-

hexyl, and the like, unless otherwise indicated.

5 "Alkenyl", alone or in combination, refers to a straight-chain or branched-chain hydrocarbon radical having one more double bonds, preferably 1-2 double bonds and more preferably one double bond, and containing a designated number of carbon atoms. For example, C₂-C₉ alkenyl is a straight or branched hydrocarbon chain containing 2 to 9 carbon atoms having at least one double bond, and includes but is not limited to ethenyl, propenyl, iso-propenyl, butenyl, 2-methylpropenyl, 10 iso-butenyl, tert-butenyl, 1,4-butadienyl, n-pentenyl, n-hexenyl, and the like, unless otherwise indicated.

15 "Alkynyl", alone or in combination, refers to a straight-chain or branched-chain hydrocarbon radical having one or more triple bonds, preferably 1-2 triple bonds and more preferably one triple bond, and containing a designated number of carbon atoms. For example, C₂-C₉ alkynyl is a straight or branched hydrocarbon chain containing 2 to 9 carbon atoms having at least one triple bond, and includes but is not limited to ethynyl, propynyl (propargyl), butynyl, and the like, unless 20 otherwise indicated.

25 "Alkanoyl", alone or in combination, means a radical of the type "R-C(O)-" wherein "R" is an alkyl radical as defined above and "-C(O)-" is a carbonyl radical. Examples of such alkanoyl radicals include without limitation acetyl, trifluoroacetyl, hydroxyacetyl, propanoyl, butyryl, pentanoyl, and the like.

30 "Alkoxy", alone or in combination, refers to a radical of the type "-OR" wherein "R" is alkyl radical as defined above and "O" is an oxygen atom. Preferably, R is a branched or unbranched saturated hydrocarbon chain containing 1 to 9 carbon atoms. Examples of such alkoxy radicals include without limitation methoxy, ethoxy, n-propoxy, isopropoxy, n-


butoxy, iso-butoxy, sec-butoxy, tert-butoxy, and the like, unless otherwise indicated.

It should be kept in mind that, throughout this application, "R" or "R_n", where n is a number, is used to designate various alkyl (and other) substituents. As indicated throughout, these R groups are independently selected. Thus, for example, the fact that R₁ may be a branched alkyl in one context does not require that R₁ be the same branched alkyl, and does not prohibit that R₁ be, for example, a straight chain alkenyl, in another context in the same molecule. It is intended that all "R_n" are selected independently of all other "R_n", whether or not the term "independently selected" is used or is inadvertently omitted.

"Aryl" or "aromatic" refers to an aromatic carbocyclic or heterocyclic moiety having one or more closed rings which is optionally substituted with one or more substituents selected from alkyl, alkoxy, halogen, hydroxy, amino, azido, nitro, cyano, haloalkyl, carboxy, alkoxycarbonyl, cycloalkyl, heterocyclo, alkanoylamino, amido, amidino, alkoxycarbonylamino, N-alkylamidino, alkylamino, dialkylamino, N-alkylamido, N,N-dialkylamido, aralkoxycarbonylamino, alkylthio, alkylsulfinyl, alkylsulfonyl, and the like. Any substituents attached to an aryl moiety in the compounds of the present invention may be configured in the ortho-, meta-, or para- orientations, with the para- orientation being preferred. Examples of preferred aryl radicals include, without limitation, phenyl, p-tolyl, 4-methoxyphenyl, 4-(tert-butoxy)phenyl, 3-methyl-4-methoxyphenyl, 4-CF₃-phenyl, 4-fluorophenyl, 4-chlorophenyl, 3-nitrophenyl, 3-aminophenyl, 3-acetamidophenyl, 4-acetamidophenyl, 2-methyl-3-acetamidophenyl, 2-methyl-3-aminophenyl, 2-amino-3-methylphenyl, 2,4-dimethyl-3-aminophenyl, 4-hydroxyphenyl, 3-

methyl-4-hydroxyphenyl, 4-(4-methoxyphenyl)phenyl, naphthyl, 3-amino-1-naphthyl, 2-methyl-3-amino-1-naphthyl, 6-amino-2-naphthyl, piperazinyphenyl, anthracenyl, phenanthracenyl, biphenyl, pyrenyl, and the like.


5 Other examples of typical aryl moieties included in the scope of the present invention may include, but are not limited to, the following:



10 It should be kept in mind that, throughout this application, "Ar" or "Ar_n", where n is a number, is used to designate various cyclic (and other) substituents. As indicated throughout, these Ar groups are independently selected. Thus, for example, the fact that Ar₂ may be phenyl
 15 in one context does not require that Ar₂ be phenyl, nor prohibit that Ar₂ be, for example, pyridyl, in another context in the same molecule. It is intended that all "Ar_n" are selected independently of all other "Ar_n", whether or not the

term "independently selected" is used or is inadvertently omitted.

5 "Bridged ring" or "bridged ring moiety", alone or in combination, refers to a carbocyclic or heterocyclic moiety where two or more atoms are shared between two or more ring structures, where any such shared atom is C, N, S, or other heteroatom arranged in a chemically reasonable substitution pattern and which is optionally substituted as defined herein with respect to the definition of aryl. Alternatively, a
10 "bridged" compound also refers to a carbocyclic or heterocyclic ring structure where one atom at any position of a primary ring is bonded to a second atom on the primary ring through either a chemical bond or atom(s) other than a bond which do not comprise a part of the primary ring structure.
15 The first and second atom may or may not be adjacent to one another in the primary ring. Illustrated below are specific nonlimiting examples of bridged ring structures contemplated by the present invention:



The present invention also contemplates other carbocyclic or heterocyclic bridged ring structures, including bridged rings wherein the bridging atoms are C or heteroatom(s) arranged in chemically reasonable substitution patterns, which are not described herein.

"Heteroaryl", alone or in combination, refers to an aromatic heterocyclic moiety having one or more closed rings with one or more heteroatoms (for example, sulfur, nitrogen or oxygen) in at least one of the rings and which is optionally substituted as defined herein with respect to the definition of aryl. Examples include, without limitation, pyrrole, thiophene, pyridine and isoxazole.

"Carbocycle" or "carbocyclic", alone or in combination, refers to a saturated or partially saturated hydrocarbon, cyclic moiety having one or more closed rings that is/are


alicyclic, aromatic, fused and/or bridged, and which is optionally substituted as defined herein with respect to the definition of aryl. "Heterocycle" or "heterocyclic", alone or in combination, refers to a saturated or partially unsaturated, cyclic moiety having one or more closed rings that is/are alicyclic, aromatic, fused and/or bridged, with one or more heteroatoms (for example, sulfur, nitrogen or oxygen) in at least one of the rings and which may or may not include carbon and which is optionally substituted as defined herein with respect to the definition of aryl. "Heterocycle" is intended to include sulfone and sulfoxide derivatives of sulfur ring members and N-oxides of tertiary nitrogen ring members, and benzofused ring systems.

Examples of preferred carbocyclic and heterocyclic moieties include, without limitation, cyclopentyl, cyclohexyl, cycloheptyl, phenyl, benzyl, naphthyl, indenyl, azulenyl, fluorenyl, anthracenyl, indolyl, isoindolyl, indolinyl, benzofuranyl, benzothiophenyl, indazolyl, benzimidazolyl, benzthiazolyl, tetrahydrofuranyl, tetrahydropyranyl, pyridyl, pyrrolyl, pyrrolidinyl, pyridinyl, pyrimidinyl, purinyl, quinolinyl, isoquinolinyl, tetrahydroquinolinyl, quinolizinyl, furyl, thiophenyl, imidazolyl, oxazolyl, benzoxazolyl, thiazolyl, isoxazolyl, isotriazolyl, oxadiazolyl, triazolyl, thiadiazolyl, pyridazinyl, pyrimidinyl, pyrazinyl, triazinyl, trithianyl, indolizinyl, pyrazolyl, pyrazolinyl, pyrazolidinyl, thienyl, tetrahydroisoquinolinyl, cinnolinyl, phthalazinyl, quinazolinyl, quinoxalinyl, naphthyridinyl, pteridinyl, carbazolyl, acridinyl, phenazinyl, phenothiazinyl, phenoxazinyl, and adamantyl.

As one skilled in the art will appreciate, such heterocyclic moieties may exist in several isomeric forms, all of which are encompassed by the present invention. For

example, a 1,3,5-triazine moiety is isomeric to a 1,2,4-triazine group. Such positional isomers are to be considered within the scope of the present invention. Likewise, the heterocyclic or heteroaryl groups can be bonded to other moieties in the compounds of the present invention. The point(s) of attachment to these other moieties is not to be construed as limiting on the scope of the invention. Thus, by way of example, a pyridyl moiety may be bound to other groups through the 2-, 3-, or 4-position of the pyridyl group. All such configurations are to be construed as within the scope of the present invention.

Examples of heterocyclic or heteroaryl moieties included in the scope of the present invention may include, but are not limited to, the following:



"Derivative" refers to a substance produced from another substance either directly or by modification or partial substitution. In particular, a "derivative" of a compound is a compound obtainable from the original compound by a simple chemical process.

"Activated derivative" of a compound means a reactive form of the original compound which renders the compound active in a desired chemical reaction, in which the original compound is only moderately reactive or non-reactive. Activation is achieved by formation of a derivative or a chemical grouping within the molecule with a higher free

energy content than that of the original compound, which renders the activated form more susceptible to react with another reagent. One skilled in the art would be able to identify various activated derivatives useful in preparing the present inventive compounds. Particularly preferred activated derivatives can include, for example, halo, a lower acyloxy group, a carbodiimide group such as 1-ethyl-3-(3-dimethylaminopropyl) carbodiimide (EDAC), an isobutyrate group, an amino acid anhydride protected amino acid, N-carboxyanhydrides (NCA's), a triazole, a mixed anhydride (see, for example, G. E. Krejcarek and K. L. Tucker, Biochem., Biophys. Res. Commun. 1977, 581, the entire contents of which are hereby incorporated by reference), such as a coupling reagent with dicyclohexyl dicarbodiimide (DCC), acid halides, anhydrides, acid chlorides, acid hydrides, activated esters, nitrenes, isothiocyanates, and acyl cyanides or anhydrides (see Tetrahedron Letters, Volume 18, (1973), pp. 1595-1598, the entire contents of which are hereby incorporated by reference).

"Effective amount" refers to the amount required to produce the desired effect. "Therapeutically effective amount" refers to the amount required to effect a desired therapeutic activity as defined herein, such as stimulation of neuronal regeneration, treating neurological disorders, treating alopecia and related hair loss disorders, treating vision disorders and/or improving vision, treating memory impairment and/or enhancing memory performance, and treating sensorineural-hearing loss. A therapeutically effective amount can be determined according to assays known to one of ordinary skill in the art, such as the MPTP assay, described herein below.

"Halo" refers to at least one fluoro, chloro, bromo or

iodo moiety.


"Isosteres" refer to elements, molecules, or ions having different molecular formulae but exhibiting similar or identical physical properties. For example, tetrazole is an isostere of carboxylic acid because it mimics the properties of carboxylic acid even though they both have very different molecular formulae. Tetrazole is one of many possible isosteric replacements for carboxylic acid. Similarly, carboxylic ester isosteres mimic the properties of an ester even though they have very different molecular formulae. Ideally, isosteric compounds should be isomorphic and able to co-crystallize. Among the other physical properties that isosteric compounds can share are boiling point, density, viscosity and thermal conductivity. The term "isosteres" encompass "bioisosteres".

"Bioisosteres" are isosteres which, in addition to their physical similarities, share some common biological properties. Typically, bioisosteres interact with the same recognition site or produce broadly similar biological effects.

"Carboxylic acid isosteres" and "carboxylic ester isosteres" include without limitation direct derivatives such as hydroxamic acids, acyl-cyanamides and acylsulfonamides; planar acidic heterocycles such as tetrazoles, mercaptoazoles, sulfinylazoles, sulfonylazoles, isoxazoles, isothiazoles, hydroxythiadiazoles and hydroxychromes; and nonplanar sulfur- or phosphorus-derived acidic functions such as phosphinates, phosphonates, phosphonamides, sulphonates, sulphonamides, and acylsulphonamides. The terms "carboxylic acid isostere" and "carboxylic ester isostere" are well known to a person of ordinary skill in the art, as evidenced by the teaching of Thornber et al., "Isosterism and molecular modification in

drug design", *Chem. Rev.* 8(4), p. 563-565 and King, "Biosisosteres, Conformational Restriction, and Pro-drugs-Case History: An Example of a Conformational Restriction Approach", *Med. Chem. Principle Practice*, 1994, P. 206-109, the contents of which are incorporated herein by reference in their entirety. Examples of these types of isosteres include, without limitation:

-COOH, -SO₃H, -SO₂HNR₃, -PO₂(R₃)₂, -CN, -PO₃(R₃)₂, -OR₃, -SR₃,
 -NHCOR₃, -N(R₃)₂, -CON(R₃)₂, -CONH(O)R₃, -CONHNHSO₂R₃, -COHNSO₂R₃,
 -CONR₃CN,



In addition, carboxylic acid isosteres can include 5-7

membered carbocycles or heterocycles containing any combination of CH₂, O, S, or N in any chemically stable oxidation state, where any of the atoms of said ring structure are optionally substituted in one or more positions.

5 The present invention contemplates that when chemical substituent(s) are added to a carboxylic acid or ester isostere, the inventive compound retains the properties of a carboxylic acid or ester isostere.

10 "Protecting group" generally refers to groups well known in the art which are used to prevent selected reactive groups, such as carboxy, amino, hydroxy, mercapto and the like, from undergoing undesired reactions, such as nucleophilic, electrophilic, oxidation, reduction, and the like. Preferred protecting groups are indicated herein where appropriate.

15 Examples of amino protecting groups include, but are not limited to, aralkyl, substituted aralkyl, cycloalkenylalkyl and substituted cycloalkenyl alkyl, allyl, substituted allyl, acyl, alkoxycarbonyl, aralkoxycarbonyl, silyl, and the like. A mixture of protecting groups can be used to protect the same

20 amino group, such as a primary amino group can be protected by both an aralkyl group and an aralkoxycarbonyl group. Amino protecting groups can also form a heterocyclic ring with the nitrogen to which they are attached, for example, 1,2-bis(methylene)benzene, phthalimidyl, succinimidyl, maleimidyl,

25 and the like and where these heterocyclic groups can further include adjoining aryl and cycloalkyl mono-, di-, or tri-substituted. Amino groups may also be protected against undesired reactions, such as oxidation, through the formation of an addition salt, such as hydrochloride, toluenesulfonic acid, trifluoroacetic acid, and the like. Many of the amino

30 protecting groups are also suitable for protecting carboxy, hydroxy, and mercapto groups. Alkyl groups are also suitable

groups for protecting hydroxy and mercapto groups.

"Low molecular weight, small molecule compounds" include, without limitation, molecules which are smaller in size, molecular weight, or both in relation to the compounds
5 Rapamycin, Cyclosporin, and FK506. Preferably, such compounds have a molecular weight no more than about 800 daltons; more preferably, no more than about 650 daltons; most preferably, no more than about 500 daltons; even more preferably, no more than about 450 daltons; more preferably, no more than about
10 400 daltons; more preferably, no more than about 350 daltons, and more preferably, no more than about 300 daltons.

"Pharmaceutically acceptable carrier" as used herein refers to any carrier, diluent, excipient, suspending agent, lubricating agent, adjuvant, vehicle, delivery system,
15 emulsifier, disintegrant, absorbent, preservative, surfactant, colorant, flavorant, or sweetener. For these purposes, the compounds of the present invention may be administered orally, parenterally, by inhalation spray, topically, rectally, nasally, buccally, vaginally, or via an implanted reservoir in
20 dosage formulations containing conventional non-toxic pharmaceutically-acceptable carriers, adjuvants, and vehicles. The term parenteral as used herein includes subcutaneous, intravenous, intramuscular, intraperitoneal, intrathecal, intraventricular, intrasternal, and intracranial injection or
25 infusion techniques.

"Pharmaceutically acceptable salt", as used herein, refers to an organic or inorganic salt of the inventive compounds which possesses the desired pharmacological activity and which is neither biologically nor otherwise undesirable.
30 Such salts can be acid or basic addition salts, depending on the nature of the inventive compound to be used.

In the case of an acidic moiety in an inventive compound,

a salt may be formed by treatment of the inventive compound with a basic compound, particularly an inorganic base. Preferred inorganic salts are those formed with alkali and alkaline earth metals such as lithium, sodium, potassium, barium and calcium. Preferred organic base salts include, for example, ammonium, dibenzylammonium, benzylammonium, 2-hydroxyethylammonium, bis(2-hydroxyethyl)ammonium, phenylethylbenzylamine, dibenzyl-ethylenediamine, and the like salts. Other salts of acidic moieties may include, for example, those salts formed with procaine, quinine and N-methylglucosamine, plus salts formed with basic amino acids such as glycine, ornithine, histidine, phenylglycine, lysine and arginine. Other suitable base salts, esters, or solvates include magnesium salts; salts with organic bases, such as dicyclohexylamine salts; and N-methyl-D-glucamine. An especially preferred salt is a sodium or potassium salt of an inventive compound. Free acids can be additionally preferred.

With respect to basic moieties, a salt is formed by the treatment of the desired inventive compound with an acidic compound, particularly an inorganic acid. Preferred inorganic salts of this type may include, for example, the hydrochloric, hydrobromic, hydroiodic, sulfuric, phosphoric or the like salts. Preferred organic salts of this type, may include, for example, salts formed with formic, acetic, succinic, citric, lactic, maleic, fumaric, palmitic, cholic, pamoic, mucic, d-glutamic, d-camphoric, glutaric, glycolic, phthalic, tartaric, lauric, stearic, salicylic, methanesulfonic, benzenesulfonic, para-toluenesulfonic, sorbic, puric, benzoic, cinnamic and the like organic acids. Other suitable acids are adipate, alginate, aspartate, benzenesulfonate, bisulfate, butyrate, camphorsulfonate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, glucoheptanoate,

glycerophosphate, hemisulfate, heptanoate, hexanoate, 2-hydroxyethanesulfonate, methanesulfonate, naphthylate, 2-naphthalenesulfonate, nicotinate, oxalate, thiocyanate, tosylate and undecanoate. An especially preferred salt of this type is a hydrochloride or sulfate salt of the desired inventive compound. Also, the basic nitrogen-containing groups can be quarternized with such agents as: 1) lower alkyl halides, such as methyl, ethyl, propyl, and butyl chloride, bromides and iodides; 2) dialkyl sulfates like dimethyl, diethyl, dibutyl and diamyl sulfates; 3) long chain alkyls such as decyl, lauryl, myristyl and stearyl substituted with one or more halide such as chloride, bromide and iodide; and 4) aralkyl halides like benzyl and phenethyl bromide and others. Free bases can be additionally preferred.

"Isomers" refer to compounds having the same number and kind of atoms (same molecular formula), and hence the same molecular weight, but differing in respect to the arrangement or configuration of the atoms.

"Stereoisomers" refer to isomers that differ only in the arrangement of the atoms in space. "Diastereoisomers" are stereoisomers which are not mirror images of each other.


"Optical isomers" refer to either of two kinds of stereoisomers. One kind is represented by mirror-image structures called enantiomers, which result from the presence of one or more asymmetric carbon atoms in the compound (glyceraldehyde, lactic acid, sugars, tartaric acid, amino acids). The other kind is exemplified by diastereoisomers, which are not mirror images. These occur in compounds having two or more asymmetric carbon atoms; thus, such compounds have 2_n optical isomers, where n is the number of asymmetric carbon atoms.

"Enantiomers" refer to a pair of stereoisomers that are

non-superimposable mirror images of each other.

"Racemic mixture" refers to a mixture containing equal parts of individual enantiomers. "Non-racemic mixture" is a mixture containing unequal parts of individual enantiomers or stereoisomers.

Further, as used throughout the teaching of the invention, a designation of:



wherein W or Y is H₂, or similar designations, is meant to denote that two hydrogen atoms are attached to the noted carbon and that the bonds to each hydrogen are single bonds.

"Alopecia" refers to deficient hair growth and partial or complete loss of hair, including without limitation androgenic alopecia (male pattern baldness), toxic alopecia, alopecia senilis, alopecia areata, alopecia pelada and trichotillomania. Alopecia results when the pilar cycle is disturbed. The most frequent phenomenon is a shortening of the hair growth or anagen phase due to cessation of cell proliferation. This results in an early onset of the catagen phase, and consequently a large number of hairs in the telogen phase during which the follicles are detached from the dermal papillae, and the hairs fall out. Alopecia has a number of etiologies, including genetic factors, aging, local and systemic diseases, febrile conditions, mental stresses, hormonal problems, and secondary effects of drugs.

"Animal" refers to a living organism having sensation and the power of voluntary movement, and which requires for its existence oxygen and organic food. Examples include, without limitation, an animal such as a member of the human, equine,

porcine, bovine, murine, canine, or feline species. A preferred animal is mammal. In the case of a human, an "animal" may also be referred to as a "patient".

5 "Disease" refers to any deviation from or interruption of the normal structure or function of any part, organ, or system (or combinations) of the body that is manifested by a characteristic set of symptoms and signs and whose etiology, pathology, and prognosis may be known or unknown. Dorland's Illustrated Medical Dictionary, (W.B. Saunders Co. 27th ed. 10 1988).

"Disorder" refers to any derangement or abnormality of function; a morbid physical or mental state. Dorland's Illustrated Medical Dictionary, (W.B. Saunders Co. 27th ed. 15 1988).

15 "Enhancing memory performance" refers to improving or increasing the mental faculty by which to register, retain or recall past experiences, knowledge, ideas, sensations, thoughts, or impressions.

"Eye" refers to the anatomical structure responsible for vision in humans and other animals, and encompasses the following anatomical structures, without limitation: lens, vitreous body, ciliary body, posterior chamber, anterior chamber, pupil, cornea, iris, canal of Schlemm, zonules of Zinn, limbus, conjunctiva, choroid, retina, central vessels of 20 the retina, optic nerve, fovea centralis, macula lutea, and sclera. 25

"Memory impairment" refers to a diminished mental registration, retention or recall of past experiences, knowledge, ideas, sensations, thoughts or impressions. Memory 30 impairment may affect short and long-term information retention, facility with spatial relationships, memory (rehearsal) strategies, and verbal retrieval and production.

Common causes of memory impairment are age, severe head trauma, brain anoxia or ischemia, alcoholic-nutritional diseases, drug intoxications and neurodegenerative diseases. For example, memory impairment is a common feature of neurodegenerative diseases such as Alzheimer's disease and senile dementia of the Alzheimer type. Memory impairment also occurs with other kinds of dementia such as multi-infarct dementia, a senile dementia caused by cerebrovascular deficiency, and the Lewy-body variant of Alzheimer's disease with or without association with Parkinson's disease. Creutzfeldt-Jakob disease is a rare dementia with which memory impairment is associated. It is a spongiform encephalopathy caused by the prion protein; it may be transmitted from other sufferers or may arise from gene mutations. Loss of memory is also a common feature of brain-damaged patients. Brain damage may occur, for example, after a classical stroke or as a result of an anaesthetic accident, head trauma, hypoglycemia, carbon monoxide poisoning, lithium intoxication, vitamin (B₁, thiamine and B₁₂) deficiency, or excessive alcohol use. Korsakoff's amnesic psychosis is a rare disorder characterized by profound memory loss and confabulation, whereby the patient invents stories to conceal his or her memory loss. It is frequently associated with excessive alcohol intake. Memory impairment may furthermore be age-associated; the ability to recall information such as names, places and words seems to decrease with increasing age. Transient memory loss may also occur in patients, suffering from a major depressive disorder, after electro-convulsive therapy.

"Neopsic factors" or "neopsics" refers to compounds useful in treating vision loss, preventing vision degeneration, or promoting vision regeneration.

"Neopsis" refers to the process of treating vision loss,

preventing vision degeneration, or promoting vision regeneration.

"Neurotrophic" includes without limitation the ability to stimulate neuronal regeneration or growth, and/or the ability to prevent or treat neurodegeneration. Assays for determining the neurotrophic abilities of a compound are well known to those of ordinary skill in the art. Specific, non-limiting examples of well known assays include MPTP and chick DRG assays. Preferably, neurotrophic compounds exhibit an MPTP Assay value which is greater than about 3% recovery of TH-stained dopaminergic neurons; more preferably, greater than about 10% recovery of TH-stained dopaminergic neurons; more preferably, greater than about 20% recovery of TH-stained dopaminergic neurons; more preferably, greater than about 35% recovery of TH-stained dopaminergic neurons; most preferably, greater than about 50% recovery of TH-stained dopaminergic neurons.

"Non-immunosuppressive" as used herein refers to the inability of the compounds of the present invention to trigger an immune response when compared to a control such as FK506 or cyclosporin A. Assays for determining immunosuppression are well known to those of ordinary skill in the art. Specific, non-limiting examples of well known assays include PMA and OKT3 wherein mitogens are used to stimulate proliferation of human peripheral blood lymphocytes (PBC) and the compounds are evaluated on their ability to inhibit such proliferation.

"Ophthalmological" refers to anything about or concerning the eye, without limitation, and is used interchangeably with "ocular", "ophthalmic", "ophthalmologic", and other such terms, without limitation.

"Pilar cycle" refers to the life cycle of hair follicles, and includes three phases:

- (1) the anagen phase, the period of active hair growth which, insofar as scalp hair is concerned, lasts about three to five years;
- (2) the catagen phase, the period when growth stops and the follicle atrophies which, insofar as scalp hair is concerned, lasts about one to two weeks; and
- (3) the telogen phase, the rest period when hair progressively separates and finally falls out which, insofar as scalp hair is concerned, lasts about three to four months.

Normally 80 to 90 percent of the follicles are in the anagen phase, less than 1 percent being in the catagen phase, and the rest being in the telogen phase. In the telogen phase, hair is uniform in diameter with a slightly bulbous, non-pigmented root. By contrast, in the anagen phase, hair has a large colored bulb at its root.

"Promoting hair growth" refers to maintaining, inducing, stimulating, accelerating, or revitalizing the germination of hair.

"Promoting vision regeneration" refers to maintaining, improving, stimulating or accelerating recovery of, or revitalizing one or more components of the visual system in a manner which improves or enhances vision, either in the presence or absence of any ophthalmologic disorder, disease, or injury.

"Preventing neurodegeneration" as used herein includes the ability to inhibit or prevent neurodegeneration in patients newly diagnosed as having a neurodegenerative disease, or at risk of developing a new degenerative disease and for inhibiting or preventing further neurodegeneration in patients who are already suffering from or have symptoms of a neurodegenerative disease when the compounds are given

concurrently.

"Preventing vision degeneration" as used herein includes the ability to prevent degeneration of vision in patients newly diagnosed as having a degenerative disease affecting vision, or at risk of developing a new degenerative disease affecting vision, and for preventing further degeneration of vision in patients who are already suffering from or have symptoms of a degenerative disease affecting vision.

"Treating" or "treatment" as used herein refers to:

(i) preventing a disease, disorder, or condition from occurring in an animal which may be predisposed to the disease, disorder, and/or condition but has not yet been diagnosed as having it;

(ii) inhibiting the disease, disorder, or condition, i.e., arresting its development; and/or

(iii) relieving the disease, disorder, or condition, i.e., causing regression of the disease, disorder, and/or condition.

"Treating alopecia" refers to:

(i) preventing alopecia in an animal which may be predisposed to alopecia; and/or

(ii) inhibiting, retarding, or reducing alopecia; and/or

(iii) promoting hair growth; and/or

(iv) prolonging the anagen phase of the hair cycle; and/or

(v) converting vellus hair to growth as terminal hair. Terminal hair is coarse, pigmented, long hair in which the bulb of the hair follicle is seated deep in the dermis. Vellus hair, on the other hand, is fine, thin, non-pigmented short hair in which the hair bulb is located superficially in the dermis. As alopecia progresses, the hairs change from the terminal to the vellus type.

"Treating memory impairment" refers to:

(i) preventing memory impairment from occurring in an animal which may be predisposed to memory impairment but has not yet been diagnosed as having it;

5 (ii) inhibiting memory impairment, i.e., arresting its development;

(iii) relieving memory impairment, i.e., causing its regression; and/or

(iv) enhancing memory.

10 "Enhancing memory performance" refers to improving or increasing the mental faculty by which to register, retain or recall past experiences, knowledge, ideas, sensations, thoughts or impressions.

15 "Vision" refers to the ability of humans and other animals to process images, and is used interchangeably with "sight", "seeing", and other such terms, without limitation.

20 "Vision disorder" refers to any disorder that affects or involves vision, including without limitation visual impairment, orbital disorders, disorders of the lacrimal apparatus, disorders of the eyelids, disorders of the conjunctiva, disorders of the cornea, cataracts, disorders of the uveal tract, disorders of the optic nerve or visual pathways, free radical induced eye disorders and diseases, immunologically-mediated eye disorders and diseases, eye injuries, and symptoms and complications of eye disease, eye disorder, or eye injury.

25 "Visual impairment" refers to any dysfunction in vision including, without limitation, disturbances or diminution in vision (e.g., binocular, central, peripheral, scotopic),
30 visual acuity for objects near and for, visual field, ocular motility, color perception, adaptation to light and dark, accommodation, refraction, and lacrimation. See *Physicians'*

Desk Reference (PDR) for Ophthalmology, 16th Edition, 6:47 (1988).

"Visual system" includes the eyes, the extraocular muscles which control eye position in the bony orbit (eye socket), the optic and other nerves that connect the eyes to the brain, and those areas of the brain that are in neural communication with the eyes.

Unless the context clearly dictates otherwise, the definitions of singular terms may be extrapolated to apply to their plural counterparts as they appear in the application; likewise, the definitions of plural terms may be extrapolated to apply to their singular counterparts as they appear in the application.

COMPOUNDS OF THE PRESENT INVENTION

The present invention relates to the surprising discovery that the inventive N-substituted cyclic aza derivative compounds are neurotrophic, are able to treat alopecia, are able to treat vision and memory disorders, and are able to treat sensorineural hearing loss. Accordingly, a novel class of cyclic aza derivative compounds is provided. These compounds may be polycyclic. Preferably, the compounds are low molecular weight, small molecule, neurotrophic, and/or N,N'-disubstituted. A preferred feature of the compounds of the present invention is that they do not exert any significant immunosuppressive activity.

The neurotrophic compounds of this invention can be periodically administered to a patient undergoing treatment for neurological disorders or for other reasons in which it is desirable to stimulate neuronal regeneration and growth, such as in various peripheral neuropathies and neurological disorders relating to neurodegeneration. The compounds of this invention can also be administered to mammals other than

humans for treatment of various mammalian neurological disorders.


In one embodiment, the compound of the present invention is an N-diketo cyclic aza derivative compound or a pharmaceutically acceptable salt, ester, or solvate thereof. In another embodiment, the compound of the present invention is an N-sulfonyl cyclic aza derivative compound, or a pharmaceutically acceptable salt, ester, or solvate thereof. In a further embodiment, the compound of the present invention is a tertiary N-aminocarbonyl cyclic aza compound, or a pharmaceutically acceptable salt, ester, or solvate thereof. In yet another embodiment, the compound of the present invention is a secondary N-aminocarbonyl cyclic aza compound, or a pharmaceutically acceptable salt, ester, or solvate thereof.

Possible substituents of any alkyl, alkenyl, aryl, heteroaryl, carbocycle or heterocycle group of the compounds of the present invention include, without limitation, C₁-C₉ straight or branched chain alkyl, C₂-C₉ straight or branched chain alkenyl, C₁-C₉ alkoxy, C₂-C₉ alkenyloxy, phenoxy, benzyloxy, C₃-C₈ cycloalkyl, C₅-C₇ cycloalkenyl, hydroxy, carboxy, carbonyl, amino, amido, cyano, isocyano, nitro, nitroso, nitrilo, isonitrilo, imino, azo, diazo, sulfonyl, sulfoxy, thio, thiocarbonyl, thiocyano, formanilido, thioformamido, sulfhydryl, halo, haloalkyl, trifluoromethyl, and carbocyclic and heterocyclic moieties. Carbocyclic moieties include alicyclic and aromatic structures.

Specific embodiments of the present inventive compounds are presented below in Tables I, II, and III. The present invention contemplates employing the compounds of Tables I, II, and III for use in compositions and methods to treat alopecia and promote hair growth in an animal, for use in

compositions and methods to treat a vision disorder, improve vision, treat memory impairment, and enhance memory performance, and for use in compositions and methods to treat a sensorineural hearing loss in an animal, and all the other uses suggested in this specification.

Preferred compounds of formula I are present below in TABLE I.



(I)

TABLE I

No	n	X	R ₁	R ₂
1	1	O	5-Phenylpentanoyl	1,1-Dimethylpropyl
2	1	O	3-Phenylpropanoyl	1,1-Dimethylpropyl
3	1	O	5-(3-Pyridyl)pent-4-ynoyl	1,1-Dimethylpropyl
4	1	O	5-(Cyano)pent-4-ynoyl	1,1-Dimethylpropyl
5	1	O	4-Phenylbutanoyl	1,1-Dimethylpropyl
6	1	O	6-Phenylhexanoyl	1,1-Dimethylpropyl
7	1	O	5-(3-Pyridyl)pentanoyl	1,1-Dimethylpropyl
8	1	O	3-Phenylpropyl ester	1,1-Dimethylpropyl
9	1	O	3-(3-Pyridyl)propyl ester	1,1-Dimethylpropyl
10	1	O	4-Phenylbutyl ester	1,1-Dimethylpropyl
11	1	O	2-Phenylethyl ester	1,1-Dimethylpropyl
12	2	O	6-Phenylhexanoyl	1,1-Dimethylpropyl
13	2	O	6-(3-Pyridyl)hexanoyl	1,1-Dimethylpropyl
14	2	O	3-Phenylpropyl ester	1,1-Dimethylpropyl
15	2	O	4-Phenylbutyl ester	1,1-Dimethylpropyl
16	2	O	5-Phenylpentyl ester	1,1-Dimethylpropyl
17	2	O	4-(3-Pyridyl)butyl ester	1,1-Dimethylpropyl
18	2	O	5-Phenylpentanoyl	1,1-Dimethylpropyl
19	1	O	COOH	3,4,5-trimethylphenyl
20	2	O	COOH	3,4,5-trimethylphenyl
21	1	O	COOH	tert-butyl
22	3	O	COOH	tert-butyl
23	1	O	COOH	cyclopentyl
24	2	O	COOH	cyclopentyl
25	3	O	COOH	cyclopentyl
26	1	O	COOH	cyclohexyl

	No	n	X	R ₁	R ₂
5	27	2	O	COOH	cyclohexyl
	28	3	O	COOH	cyclohexyl
	29	1	O	COOH	cycloheptyl
	30	2	O	COOH	cycloheptyl
	31	3	O	COOH	cycloheptyl
10	32	1	O	COOH	2-thienyl
	33	2	O	COOH	2-thienyl
	34	3	O	COOH	2-thienyl
	35	1	O	COOH	2-furyl
	36	2	O	COOH	3-furyl
15	37	3	O	COOH	4-furyl
	38	3	O	COOH	phenyl
	39	1	O	COOH	1,1-dimethylpentyl
	40	2	O	COOH	1,1-dimethylhexyl
	41	3	O	COOH	ethyl
20	42	1	O	SO ₃ H	1,1-dimethylpropyl
	43	1	O	CN	1,1-dimethylpropyl
	44	1	O	tetrazole	1,1-dimethylpropyl
	45	1	O	CONH ₂	1,1-dimethylpropyl
	46	2	O	CONH ₂	1,1-dimethylpropyl
25	47	1	O	COOH	α-methylbenzyl
	48	2	O	COOH	4-methylbenzyl
	49	1	O	tetrazole	benzyl
	50	1	O	SO ₃ H	α-methylbenzyl
	51	1	O	SO ₂ HNMe	benzyl
30	52	1	O	CN	α-methylbenzyl
	53	1	O	PO ₃ H ₂	4-methylbenzyl
	54	2	O	COOH	benzyl
	55	2	O	COOH	α-methylbenzyl
	56	2	O	COOH	4-methylbenzyl
35	57	2	O	COOH	cyclohexyl
	58	2	O	PO ₂ HEt	i-propyl
	59	2	O	PO ₃ HPropyl	ethyl
	60	2	O	PO ₃ (Et) ₂	methyl
	61	2	O	methyl ester	tert-butyl
40	62	1	O	ethyl ester	n-pentyl
	63	2	O	propyl ester	n-hexyl
	64	1	O	butyl ester	cyclohexyl
	65	1	O	pentyl ester	cyclopentyl
	66	1	O	hexyl ester	n-heptyl
45	67	1	O	S-Me	n-octyl
	68	1	O	S-Et	n-nonyl
	69	2	O	S-propyl	2-indolyl
	70	2	O	S-butyl	2-furyl
	71	2	O	NHCOMe	2-thiazolyl
	72	2	O	NHCOEt	2-thienyl

No	n	X	R ₁	R ₂
73	1	O	CONH(O)Me	benzyl
74	1	O	CONH(O)Et	α-methylphenyl
75	1	O	CONH(O)propyl	4-methylphenyl
76	3	O	CONHNHSO ₂ Me	benzyl
5 77	3	O	CONHNHSO ₂ Et	α-methylphenyl
78	3	O	CONHSO ₂ Me	4-methylphenyl
79	1	O	CONHNHSO ₂ Et	phenyl
80	2	O	CON(Me)CN	α-methylphenyl
81	1	O	CON(Et)CN	4-methylphenyl
10 82	1	O	COOH	1,1-dimethylpropyl
83	2	O	COOH	1,1-dimethylpropyl
84	2	O	5-(3-pyridyl)pentyl ester	1,1-dimethylpropyl
85	1	O	4-(3-pyridyl)-3-butynyl ester	1,1-dimethylpropyl
86	1	O	3-butynyl ester	1,1-dimethylpropyl
15 87	1	O	5-phenylpentyl ester	1,1-dimethylpropyl
88	1	O	4-(3-pyridyl)butyl ester	1,1-dimethylpropyl
89	1	O	3-phenylpropyl ester	1,1-dimethylpentyl
90	1	O	3-(3-pyridyl)propyl ester	1,1-dimethylpentyl
91	1	O	4-phenylbutyl ester	1,1-dimethylpentyl
20 92	1	O	2-phenylethyl ester	1,1-dimethylpropyl
93	1	O	2-phenylethanoyl	1,1-dimethylpropyl
94	2	O	5-(3-pyridyl)pentanoyl	1,1-dimethylpropyl
95	2	O	4-phenylbutanoyl	1,1-dimethylpropyl
96	1	O	4-(3-pyridyl)butanoyl	1,1-dimethylpropyl
25 97	2	S	2-phenylethyl ester	1,1-dimethylpropyl
98	2	S	3-phenylpropyl ester	1,1-dimethylpropyl
99	1	S	3-phenylpropyl ester	1,1-dimethylpropyl
100	1	S	2-phenethylester	1,1-dimethylpropyl
101	1	S	COOH	1,1-dimethylpropyl
30 102	2	S	PO ₃ H ₂	2-furyl
103	1	S	COOH	phenyl
104	2	S	COOH	3,4,5-trimethoxyphenyl

Preferred compounds of TABLE I are named as follows:

35 3-phenyl-1-propyl 1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyridazinecarboxylate,

 4-phenyl-1-n-butyl 1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyridazinecarboxylate,

40 5-phenyl-1-n-pentyl 1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyridazinecarboxylate,

4-(3-pyridyl)-1-n-butyl 1-(3,3-dimethyl-1,2-dioxopentyl)-
2-pyridazinecarboxylate,

3-phenyl-1-propyl 1-(3,3-dimethyl-1,2-dioxopentyl)-2-
pyrazinecarboxylate,

5 3-(3-pyridyl)-1-propyl 1-(3,3-dimethyl-1,2-dioxopentyl)-
2-pyrazinecarboxylate,

4-phenyl-1-n-butyl 1-(3,3-dimethyl-1,2-dioxopentyl)-2-
pyrazinecarboxylate,

10 2-phenyl-1-ethyl 1-(3,3-dimethyl-1,2-dioxopentyl)-2-
pyrazinecarboxylate,

2-[(4-phenylbutyl)carbonyl]-1-(3,3-dimethyl-1,2-
dioxopentyl)pyridazine,

2-[(2-phenylethyl)carbonyl]-1-(3,3-dimethyl-1,2-
dioxopentyl)pyridazine,

15 2-[(5-phenylpentyl)carbonyl]-1-(3,3-dimethyl-1,2-
dioxopentyl)piperazine,

2-[(5-(3-pyridyl)pentyl)carbonyl]-1-(3,3-dimethyl-1,2-
dioxopentyl)piperazine,

20 2-[(4-phenylbutyl)carbonyl]-1-(3,3-dimethyl-1,2-
dioxopentyl)piperazine,

2-[(3-phenylpropyl)carbonyl]-1-(3,3-dimethyl-1,2-
dioxopentyl)pyridazine,

2-[(5-phenylpentyl)carbonyl]-1-(3,3-dimethyl-1,2-
dioxopentyl)pyridazine, and

25 2-[(4-(3-pyridyl)butyl)carbonyl]-1-(3,3-dimethyl-1,2-
dioxopentyl)pyridazine.

Preferred compounds of formula II are present below in
TABLE II.




TABLE II

	No.	n	R ₁	R ₂
5	105	1	3-Phenylpropyl ester	benzyl
	106	2	4-Phenylbutyl ester	benzyl
	107	1	5-Phenylpentanoyl	benzyl
	108	1	COOH	benzyl
	109	1	COOH	α-methylbenzyl
10	110	1	COOH	4-methylbenzyl
	111	1	tetrazole	benzyl
	112	1	SO ₃ H	α-methylbenzyl
	113	1	SO ₂ HMe	benzyl
	114	1	CN	α-methylbenzyl
15	115	1	PO ₃ H ₂	4-methylbenzyl
	116	2	COOH	benzyl
	117	2	COOH	α-methylbenzyl
	118	2	COOH	4-methylbenzyl
	119	2	COOH	3,4,5-trimethoxyphenyl
20	120	2	COOH	cyclohexyl
	121	2	PO ₂ HEt	i-propyl
	122	2	PO ₃ HPropyl	ethyl
	123	2	PO ₃ (Et) ₂	methyl
	124	2	methyl ester	tert-butyl
25	125	2	ethyl ester	n-pentyl
	126	2	propyl ester	n-hexyl
	127	1	butyl ester	cyclohexyl
	128	1	pentyl ester	cyclopentyl
	129	1	hexyl ester	n-heptyl
30	130	1	S-Me	n-octyl
	131	1	S-Et	n-nonyl
	132	2	S-propyl	2-indolyl
	133	2	S-butyl	2-furyl
	134	2	NHCOMe	2-thiazolyl
35	135	2	NHCOEt	2-thienyl
	136	1	CONH(O)Me	benzyl
	137	1	CONH(O)Et	α-methylphenyl
	138	1	CONH(O)propyl	4-methylphenyl
	139	2	COOH	benzyl
40	140	2	COOH	α-methylphenyl
	141	2	COOH	4-methylphenyl


No.	n	R ₁	R ₂
142	3	CONHNHSO ₂ Me	benzyl
143	3	CONHNHSO ₂ Et	α-methylphenyl
144	3	CONHSO ₂ Me	4-methylphenyl
145	2	CONHNHSO ₂ Et	phenyl
146	2	CON(Me)CN	α-methylphenyl
147	2	CON(Et)CN	4-methylphenyl

Preferred compounds of TABLE II are named as follows:

4-phenyl-1-n-butyl 1-(phenylmethyl)sulfonyl-2-pyridazinecarboxylate, and

3-phenyl-1-propyl 1-(phenylmethyl)sulfonyl-2-pyrazinecarboxylate.

Preferred compounds of formula III are present below in TABLE III.



(III)

TABLE III

No.	n	R ₁	R ₄	R ₅
148	1	5-Phenylpentanoyl	cyclohexyl	cyclohexyl
149	1	COOH	cyclohexyl	methyl
150	1	COOH	cyclohexyl	ethyl
151	1	COOH	cyclohexyl	propyl
152	1	COOH	cyclohexyl	butyl

Preferred compounds of formula IV are present below in TABLE IV.




TABLE IV

	No.	n	R ₁	R ₂
5	153	1	3-phenylpropyl ester	cyclohexyl
	154	2	4-phenylbutyl ester	cyclohexyl
	155	1	5-phenylpentanoyl	cyclohexyl
	156	1	COOH	cyclohexyl
	157	1	COOH	α-methylbenzyl
10	158	1	COOH	4-methylbenzyl
	159	1	tetrazole	benzyl
	160	1	SO ₃ H	α-methylbenzyl
	161	1	SO ₂ HNMe	benzyl
	162	1	CN	α-methylbenzyl
15	163	1	PO ₃ H ₂	4-methylbenzyl
	164	2	COOH	benzyl
	165	2	COOH	α-methylbenzyl
	166	2	COOH	2-butyl
	167	2	COOH	cyclohexyl
20	168	2	PO ₂ HEt	i-propyl
	169	2	PO ₃ HPropyl	ethyl
	170	2	PO ₃ (Et) ₂	methyl
	171	2	Methyl ester	tert-butyl
	172	2	Ethyl ester	n-pentyl
25	173	2	propyl ester	n-hexyl
	174	1	butyl ester	cyclohexyl
	175	1	pentyl ester	cyclopentyl
	176	1	hexyl ester	heptyl
	177	1	SMe	n-octyl
30	178	1	SEt	n-hexyl
	179	2	S-propyl	n-hexyl
	180	2	S-butyl	n-hexyl
	181	2	NHCOMe	n-hexyl
	182	2	NHCOEt	2-thienyl
35	183	1	CONH(O)Me	benzyl
	184	1	CONH(O)Et	α-methylphenyl
	185	1	CONH(O)propyl	4-methylphenyl
	186	2	COOH	benzyl
	187	2	COOH	α-methylphenyl
	188	2	COOH	4-methylphenyl



No.	n	R ₁	R ₂
189	3	CONHNHSO ₂ Me	benzyl
190	3	CONHNHSO ₂ Et	α-methylphenyl
191	3	CONHSO ₂ Me	4-methylphenyl
192	2	CONHNHSO ₂ Et	phenyl
193	2	CON(Me)CN	α-methylphenyl
194	2	CON(Et)CN	4-methylphenyl
195	1	3-phenylpropyl ester	cyclohexyl





Preferred compounds of TABLE IV are named as follows:





4-phenyl-1-n-butyl 1-(cyclohexyl)carbamoyl-2-pyridazinecarboxylate, and






3-phenyl-1-propyl 1-(cyclohexyl)carbamoyl-2-pyrazinecarboxylate.






Representative compounds of the present invention are set forth below.





No.	Structure	Name
1		3,3-dimethyl-N-[2-(5-phenylpentanoyl)-tetrahydro-1H-1-pyrazolyl]-1,2-pentanedione
2		3,3-dimethyl-N-[2-(3-phenylpropanoyl)-tetrahydro-1H-1-pyrazolyl]-1,2-pentanedione


No.	Structure	Name
3		3,3-dimethyl-1-[2-(5-(3-pyridyl)pent-4-ynoyl)-pyrazolidinyl]-pentane-1,2-dione
4		3,3-dimethyl-1-[2-(5-(cyano)pent-4-ynoyl)-pyrazolidinyl]-pentane-1,2-dione
5		3,3-dimethyl-1-[2-(4-phenylbutanoyl)-pyrazolidinyl]-pentane-1,2-dione
6		3,3-dimethyl-1-[2-(6-phenylhexanoyl)-pyrazolidinyl]-pentane-1,2-dione

No.	Structure	Name
7		3,3-dimethyl-1-[2-(5-(3-pyridyl)pentanoyl)-pyrazolidinyl]pentane-1,2-dione
8		3-phenylpropyl 2-(3,3-dimethyl-2-oxopentanoyl)-pyrazolidinecarboxylate
9		3-(3-pyridyl)propyl 2-(3,3-dimethyl-2-oxopentanoyl)-pyrazolidinecarboxylate
10		4-phenylbutyl 2-(3,3-dimethyl-2-oxopentanoyl)-pyrazolidinecarboxylate

No.	Structure	Name
11		2-phenylethyl 2-(3,3-dimethyl-2-oxopentanoyl)-pyrazolidinecarboxylate
12		3,3-dimethyl-1-[2-(6-phenylhexanoyl)-perhydropyridazinyl]-pentane-1,2-dione
13		3,3-dimethyl-2-[2-(6-(3-pyridyl)hexanoyl)-perhydropyridazinyl]-pentane-1,2-dione
14		3-phenylpropyl 2-(3,3-dimethyl-2-oxopentanoyl)-perhydropyridazinecarboxylate
15		4-phenylbutyl 2-(3,3-dimethyl-2-oxopentanoyl)perhydropyridazinecarboxylate

No.	Structure	Name
16		5-phenylpentyl 2-(3,3-dimethyl-2-oxopentanoate)perhydropyridazinecarboxylate
17		4-(3-pyridyl)butyl 2-(3,3-dimethyl-2-oxopentanoate)perhydropyridazinecarboxylate
18		3,3-dimethyl-1-[2-(5-phenylpentanoyl)perhydropyridazinyl]pentane-1,2-dione
105		3-phenylpropyl 2-[benzylsulfonyl]pyrazolidinecarboxylate
106		4-phenylbutyl 2-[benzylsulfonyl]perhydropyridazinecarboxylate

No.	Structure	Name
107		1-(5-phenylpentanoyl)-2-(benzylsulfonyl)-tetrahydro-1H-1-pyrazole
153		3-phenylpropyl 2-(N-cyclohexylcarbamoyl)-pyrazolidinecarboxylate
154		4-phenylbutyl 2-(N-cyclohexylcarbamoyl)-perhydropyridazinecarboxylate
155		1-(5-phenylpentanoyl)-2-(N-cyclohexylcarbamoyl)tetrahydro-1H-1-pyrazole

No.	Structure	Name
148		1-(5-phenylpentanoyl)-2-(N,N-dicyclohexylcarbamoyl)tetrahydro-1H-pyrazole

Other compounds which are cyclic aza derivative compounds within the scope of the present invention are those compounds which may possess immunosuppressive, non-immunosuppressive, or other activities as long as they also are useful in treating a disease. In particular, other cyclic aza derivative compounds falling within the scope of the present invention are useful for preventing and/or treating neurological disorders, including physically damaged nerves and neurodegenerative diseases; in treating alopecia and promoting hair growth; in treating vision disorders and/or improving vision; in treating memory impairment and/or enhancing memory performance; and/or in treating sensorineural hearing loss.

The compounds of this invention may possess at least one asymmetric center and thus can be produced as mixtures of stereoisomers or as individual R- and S-stereoisomers. The individual enantiomers may be obtained by using an optically active starting material, by resolving a racemic or non-racemic mixture of an intermediate at some appropriate stage of the synthesis, or by resolving a compound of the present invention. It is understood that the individual R- and S-stereoisomers as well as mixtures (racemic and non-racemic) of stereoisomers are encompassed by this invention.

PHARMACEUTICAL COMPOSITIONS OF THE PRESENT INVENTION

The present invention also relates to a pharmaceutical composition comprising:

(i) an effective amount of a compound of the present invention, as defined above; and

(ii) a pharmaceutically acceptable carrier.

Preferably, the compound is present in an effective amount for effecting a neuronal activity.

The present invention also relates to a pharmaceutical composition comprising:

(i) an effective amount of a compound of the present invention, as defined above, for treating neurodegenerative diseases, neurological disorders, and nerve damage, or promoting nerve growth in animals; and

(ii) a pharmaceutically acceptable carrier.

The present invention also relates to a pharmaceutical composition comprising:

(i) an effective amount of a compound of the present invention, as defined above, for treating alopecia or promoting hair growth in an animal; and

(ii) a pharmaceutically acceptable carrier.

The present invention also relates to a pharmaceutical composition comprising:

(i) an effective amount of a compound of the present invention, as defined above, for treating a vision disorder, improving vision, treating memory impairment, or enhancing memory performance in an

animal; and

(ii) a pharmaceutically acceptable carrier.

The present invention also relates to a pharmaceutical composition comprising:

5 (i) an effective amount of a compound of the present invention, as defined above, for treating sensorineural hearing loss in an animal; and

(ii) a pharmaceutically acceptable carrier.

10 In another preferred embodiment, the novel compounds of the present invention can be administered in pharmaceutical compositions additionally containing other neurotrophic agents such as neurotrophic growth factor, brain derived growth factor, glial derived growth factor, ciliary neurotrophic factor, insulin growth factor and active truncated derivatives
15 thereof, acidic fibroblast growth factor, basic fibroblast growth factor, platelet-derived growth factors, neurotrophin-3 and neurotrophin 4/5. The dosage level of other neurotrophic drugs will depend upon the factors stated elsewhere herein and the neurotrophic effectiveness of the drug combination.

20 **METHODS OF THE PRESENT INVENTION**

The present invention further relates to the use of any of the compounds of the present invention, as defined above, and other compounds not specifically mentioned or described herein, in the preparation of a medicament for the treatment
25 of a disease. In particular, the compounds are used in the preparation of a medicament for treating any of the diseases enumerated herein.

The present invention also relates to the use of any of the compounds of the present invention, as defined above, and
30 other compounds not specifically mentioned or described

herein, for the treatment of a disease. In particular, the compounds are used for treating any of the diseases enumerated herein.

5 The present invention also relates to methods of using any the compounds of the present invention, as defined above, and other compounds not specifically mentioned or described herein, for treating any of the diseases enumerated herein.

10 The discussion below relating to the utility and administration of the compounds of the present invention also applies to the pharmaceutical compositions of the present invention.

METHODS FOR EFFECTING NEURONAL ACTIVITIES

15 The novel compounds of the present invention possess an excellent degree of neurotrophic activity. Accordingly, the present invention further relates to a method for effecting a neuronal activity in a mammal, comprising administering to said mammal an effective amount of a compound of the present invention, as defined above.

20 The neuronal activity that is effected by the inventive method may be selected from the group consisting of stimulation of damaged neurons, promotion of neuronal regeneration, prevention of neurodegeneration, and treatment of a neurological disorder.

25 Examples of neurological disorders that are treatable by the methods of the present invention include without limitation: trigeminal neuralgia; glossopharyngeal neuralgia; Bell's Palsy; myasthenia gravis; muscular dystrophy; amyotrophic lateral sclerosis; progressive muscular atrophy; progressive bulbar inherited muscular atrophy; herniated, 30 ruptured or prolapsed invertebrate disk syndromes; cervical

5 spondylosis; plexus disorders; thoracic outlet destruction
syndromes; peripheral neuropathies such as those caused by
lead, dapson, ticks, porphyria, or Guillain-Barré syndrome;
multiple sclerosis; stroke and ischemia associated with
stroke; neural paropathy; other neurodegenerative diseases;
motor neuron diseases; sciatic crush; sciatica; sciatic nerve
damage; peripheral neuropathy, particularly neuropathy
associated with diabetes; spinal cord injuries; facial nerve
crush; Alzheimer's disease; Huntington's disease; and
10 Parkinson's disease.

The inventive method is particularly useful for treating
a neurological disorder selected from the group consisting of
peripheral neuropathy caused by physical injury or disease
state, physical damage to the brain, physical damage to the
15 spinal cord, stroke associated with brain damage,
demyelinating disease and neurological disorder relating to
neurodegeneration.

Examples of neurological disorders relating to
neurodegeneration include Alzheimer's disease, Parkinson's
20 disease, Huntington's disease and amyotrophic lateral
sclerosis (ALS).

METHODS FOR TREATING ALOPECIA OR PROMOTING HAIR GROWTH

The present invention further relates to a method for
treating alopecia or promoting hair growth in a mammal,
25 comprising administering to the mammal an effective amount of
a compound of the present invention, as defined above.

The inventive method is particularly useful for treating
male pattern alopecia, alopecia senilis, alopecia areata,
alopecia resulting from skin lesions or tumors, alopecia
30 resulting from cancer therapy such as chemotherapy and
radiation, and alopecia resulting from systematic disorders

such as nutritional disorders and internal secretion disorders.

METHODS FOR TREATING VISION DISORDERS OR IMPROVING VISION

5 The present invention further relates to a method for treating a vision disorder, promoting vision regeneration, or improving vision in a mammal, comprising administering to the mammal an effective amount of a compound of the present invention, as defined above.

10 The inventive methods are particularly useful for treating various eye disorders including, but not limited to visual disorders, diseases, injuries, and complications, genetic disorders; disorders associated with aging or degenerative vision diseases; vision disorders correlating to physical injury to the eye, head, or other parts of the body
15 resulting from external forces; disorders resulting from environmental factors; disorders resulting from a broad range of diseases; and combinations of any of the above.

20 In particular, the compositions and methods of the present invention are useful for improving vision, or correcting, treating, or preventing visual (ocular) impairment or dysfunction of the visual system, including permanent and temporary visual impairment, without limitation. The present invention is also useful in preventing and treating
25 ophthalmologic diseases and disorders, treating damaged and injured eyes, and preventing and treating diseases, disorders, and injuries which result in vision deficiency, vision loss, or reduced capacity to see or process images, and the symptoms and complications resulting from same. The eye diseases and disorders which may be treated or prevented by the
30 compositions and methods of the present invention are not limited with regard to the cause of said diseases or

disorders. Accordingly, said compositions and methods are applicable whether the disease or disorder is caused by genetic or environmental factors, as well as any other influences. The compositions and methods of the present invention are particularly useful for eye problems or vision loss or deficiency associated with all of the following, without limitation: aging, cellular or physiological degeneration, central nervous system or neurological disorder, vascular defects, muscular defects, and exposure to adverse environmental conditions or substances.

The compositions and methods of the present invention are particularly useful in correcting, treating, or improving visual impairment, without limitation. Visual impairment in varying degrees occurs in the presence of a deviation from normal in one or more functions of the eye, including (1) visual acuity for objects at distance and near; (2) visual fields; and (3) ocular motility without diplopia. See *Physicians' Desk Reference (PDR) for Ophthalmology*, 16th Edition, 6:47 (1988). Vision is imperfect without the coordinated function of all three. *Id.*

Said compositions and methods of use are also useful in correcting, treating, or improving other ocular functions including, without limitation, color perception, adaptation to light and dark, accommodation, metamorphopsia, and binocular vision. The compositions and methods of use are particularly useful in treating, correcting, or preventing ocular disturbances including, without limitation, paresis of accommodation, iridoplegia, entropion, ectropion, epiphora, lagophthalmos, scarring, vitreous opacities, non-reactive pupil, light scattering disturbances of the cornea or other media, and permanent deformities of the orbit.

The compositions and methods of use of the present invention are also highly useful in improving vision and treating vision loss. Vision loss ranging from slight loss to absolute loss may be treated or prevented using said compositions and methods of use. Vision may be improved by the treatment of eye disorders, diseases, and injuries using the compositions and methods of the invention. However, improvements in vision using the compositions and methods of use are not so limited, and may occur in the absence of any such disorder, disease, or injury.

METHODS FOR TREATING MEMORY IMPAIRMENT OR ENHANCING MEMORY PERFORMANCE

The present invention further relates to a method for treating memory impairment or enhancing memory performance in a mammal, comprising administering to the mammal an effective amount of a compound of the present invention, as defined above.

METHODS FOR TREATING HEARING LOSS

The present invention further relates to a method for treating a sensorineural hearing loss in a mammal, comprising administering to the mammal an affective amount of the compound of the present invention, as defined above.

It is further contemplated that administration of an inventive compound will protect hair cells and spiral ganglion neurons from traumatic damage, for example damage caused by noise trauma, acute or chronic treatment with cisplatin and aminoglycoside antibiotics or from damage resulting from a lack of neurotrophic factors resulting from interruption of transport of the factors from the axon to the cell body. Such

treatment is expected to allow hair cells and/or auditory neurons to tolerate intermittent insults from either environmental noise trauma or treatment with ototoxins, and to slow down, prevent or reverse the progressive degeneration of the auditory neurons and hair cells which is responsible for hearing loss in pathological conditions such as presbycusis (age-related hearing loss), inherited sensorineural degeneration, and post-idiopathic hearing losses and to preserve the functional integrity of the inner ear. Such treatment will also support the auditory neurons for better and longer performance of cochlear implants.


METHODS FOR PREPARING INVENTIVE COMPOUNDS

The compounds of the present invention can be readily prepared by standard techniques of organic chemistry, utilizing the general synthetic pathways depicted below in Schemes I and II.

In the preparation of the compounds of the present invention, one skilled in the art will understand that one may need to protect or block various reactive functionalities on the starting compounds or intermediates while a desired reaction is carried out on other portions of the molecule. After the desired reactions are complete, or at any desired time, normally such protecting groups will be removed under conditions which will not affect the remaining portion of the molecule, for example by hydrolytic or hydrogenolytic means and the like. Such protection and deprotection steps are conventional in organic chemistry. One skilled in the art is referred to "Protective Groups in Organic Chemistry," McOmie, ed., Plenum Press, New York, New York; and "Protective Groups in Organic Synthesis," Greene, ed., John Wiley & Sons, New York, New York (1981) for the teaching of protective groups

which may be useful in the preparation of compounds of the present invention. A preferred method involves removal of a protecting group, such as removal of a benzyloxycarbonyl group by hydrogenolysis utilizing palladium on carbon in a suitable solvent system such as an alcohol, acetic acid, and the like or mixtures thereof. A t-butoxycarbonyl protecting group can be removed utilizing an inorganic or organic acid, such as HCl or trifluoroacetic acid, in a suitable solvent system, such as dioxane or methylene chloride. The resulting amino salt can be readily neutralized to yield the free amine. Carboxy protecting group, such as methyl, ethyl, benzyl, tert-butyl, 4-methoxyphenylmethyl and the like, can be removed under hydrolysis and hydrogenolysis conditions well known to those skilled in the art.

The product and intermediates may be isolated or purified using one or more standard purification techniques, including, for example, one or more of simple solvent evaporation, recrystallization, distillation, sublimation, filtration, chromatography, including thin-layer chromatography, HPLC (e.g. reverse phase HPLC), column chromatography, flash chromatography, radial chromatography, trituration, and the like.




SCHEME I

wherein, in Scheme I, n , R_3 , and R_2 are as defined elsewhere throughout the specification; R' is a straight or branched

chain alkyl group which is optionally substituted in one or more positions; and X is a halogen, wherein any of these substituents are formed in any chemically reasonable substitution pattern. It is further contemplated as within
5 the scope of the present invention that the chlorine atoms depicted in Scheme I above can be replaced with any other halogen atom.

In particular, the process of scheme I further contemplates activated derivatives of any of the compounds embodied therein, as well as the removal of a protecting group
10 from the product. One skilled in the art would be able to identify various activated derivatives useful in preparing the present inventive compounds. Particularly preferred activated derivatives can include, for example, halo, a lower acyloxy group, a carbodiimide group such as 1-ethyl-3-(3-
15 dimethylaminopropyl) carbodiimide (EDAC), an isobutyrate group, an amino acid anhydride protected amino acid, N-carboxyanhydrides (NCA's), a triazole, a mixed anhydride (see, for example, G. E. Krejcarek and K. L. Tucker, Biochem.,
20 Biophys. Res. Commun. 1977, 581, the entire contents of which are hereby incorporated by reference), such as a coupling reagent with dicyclohexyl dicarbodiimide (DCC), acid halides, anhydrides, acid chlorides, acid hydrides, activated esters, nitrenes, isothiocyanates, and acyl cyanides or anhydrides
25 (see Tetrahedron Letters, Volume 18, (1973), pp. 1595-1598, the entire contents of which are hereby incorporated by reference).



SCHEME II

wherein, in Scheme II, n , R_1 , and R_2 are as defined elsewhere throughout the specification; R' is a straight or branched chain alkyl group which is optionally substituted in one or more positions; and X is a halogen, wherein any of these substituents are formed in any chemically reasonable substitution pattern. It is further contemplated as within the scope of the present invention that the benzyl groups depicted in Scheme II above can be replaced with any R_4 group, wherein R_4 is an alkyl chain substituted with an aryl group; and that the chlorine atoms depicted in Scheme II above can be replaced with any other halogen atom.

In particular, the process of scheme II further contemplates activated derivatives of any of the compounds embodied therein, as well as the removal of a protecting group from the product. One skilled in the art would be able to identify various activated derivatives useful in preparing the present inventive compounds. Particularly preferred activated derivatives can include, for example, halo, a lower acyloxy group, a carbodiimide group such as 1-ethyl-3-(3-dimethylaminopropyl) carbodiimide (EDAC), an isobutyrate group, an amino acid anhydride protected amino acid, N-carboxyanhydrides (NCA's), a triazole, a mixed anhydride (see, for example, G. E. Krejcarek and K. L. Tucker, Biochem., Biophys. Res. Commun. 1977, 581, the entire contents of which are hereby incorporated by reference), such as a coupling reagent with dicyclohexyl dicarbodiimide (DCC), acid halides, anhydrides, acid chlorides, acid hydrides, activated esters, nitrenes, isothiocyanates, and acyl cyanides or anhydrides (see Tetrahedron Letters, Volume 18, (1973), pp. 1595-1598, the entire contents of which are hereby incorporated by reference).

ROUTE OF ADMINISTRATION

In the inventive methods, the compounds will generally be administered to a patient in the form of a pharmaceutical formulation. Such formulation preferably includes, in addition to the active agent, a physiologically acceptable carrier and/or diluent. The compounds may be administered by any means known to an ordinarily skilled artisan. For example, the compounds may be administered orally, parenterally, by inhalation spray, topically, rectally, nasally, buccally, vaginally, directly into the middle or inner ear, or via an implanted reservoir in dosage formulations containing conventional non-toxic pharmaceutically-acceptable carriers, adjuvants and vehicles. The term parenteral as used herein includes subcutaneous, intravenous, intramuscular, intraperitoneal, intrathecal, intraventricular, intrasternal, intracranial, intracerebral, intraosseous, infusion, transdermal, and transpulmonary injection or infusion routes.

For oral administration, the compounds of the present invention may be provided in any suitable dosage form known in the art. For example, the compositions may be incorporated into tablets, powders, granules, beads, chewable lozenges, capsules, liquids, aqueous suspensions or solutions, or similar dosage forms, using conventional equipment and techniques known in the art. Tablet dosage forms are preferred. Tablets may contain carriers such as lactose and corn starch, and/or lubricating agents such as magnesium stearate. Capsules may contain diluents including lactose and dried corn starch. Aqueous suspensions may contain emulsifying and suspending agents combined with the active ingredient.

When preparing dosage form incorporating the compositions of the invention, the compounds may also be blended with conventional excipients such as binders, including gelatin, pregelatinized starch, and the like; lubricants, such as hydrogenated vegetable oil, stearic acid, and the like; diluents, such as lactose, mannose, and sucrose; disintegrants, such as carboxymethylcellulose and sodium starch glycolate; suspending agents, such as povidone, polyvinyl alcohol, and the like; absorbants, such as silicon dioxide; preservatives, such as methylparaben, propylparaben, and sodium benzoate; surfactants, such as sodium lauryl sulfate, polysorbate 80, and the like; colorants such as F.D. & C. dyes and lakes; flavorants; and sweeteners.

Compositions and methods of the invention also may utilize controlled release technology. Thus, for example, the inventive compounds may be incorporated into a hydrophobic polymer matrix for controlled release over a period of days. Such controlled release films are well known to the art. Particularly preferred are transdermal delivery systems. Other examples of polymers commonly employed for this purpose that may be used in the present invention include nondegradable ethylene-vinyl acetate copolymer and degradable lactic acid-glycolic acid copolymers which may be used externally or internally. Certain hydrogels such as poly(hydroxyethylmethacrylate) or poly(vinylalcohol) also may be useful, but for shorter release cycles than the other polymer releases systems, such as those mentioned above.

The compounds of the present invention may be administered in the form of sterile injectable preparations, for example, as sterile injectable aqueous or oleaginous suspensions. These suspensions may be formulated according to techniques known in the art using suitable dispersing or

wetting agents and suspending agents. The sterile injectable preparations may also be sterile injectable solutions or suspensions in non-toxic parenterally-acceptable diluents or solvents, for example, as solutions in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as solvents or suspending mediums. For this purpose, any bland fixed oil may be employed including synthetic mono- or di-glycerides. Fatty acids such as oleic acid and its glyceride derivatives, including olive oil and castor oil, especially in their polyoxyethylated versions, are useful in the preparation of injectables. These oil solutions or suspensions may also contain long-chain alcohol diluents or dispersants.

The compounds of this invention may also be administered rectally in the form of suppositories. These compositions can be prepared by mixing the drug with a suitable non-irritating excipient which is solid at room temperature, but liquid at rectal temperature and, therefore, will melt in the rectum to release the drug. Such materials include cocoa butter, beeswax and polyethylene glycols.

The compounds of this invention may also be administered topically, especially when the conditions addressed for treatment involve areas or organs readily accessible by topical application, including neurological disorders of the eye, the skin, or the lower intestinal tract. Suitable topical formulations are readily prepared for each of these areas.

For topical application to the eye, or ophthalmic use, the compounds can be formulated as micronized suspensions in

isotonic, pH adjusted sterile saline, or, preferably, as solutions in isotonic, pH adjusted sterile saline, either with or without a preservative such as benzylalkonium chloride. Alternatively for the ophthalmic uses the compounds may be formulated in an ointment such as petrolatum.

For topical application to the skin, the compounds can be formulated in a suitable ointment containing the compound suspended or dissolved in, for example, a mixture with one or more of the following: mineral oil, liquid petrolatum, white petrolatum, propylene glycol, polyoxyethylene polyoxypropylene compound, emulsifying wax and water. Alternatively, the compounds can be formulated in a suitable lotion or cream containing the active compound suspended or dissolved in, for example, a mixture of one or more of the following: mineral oil, sorbitan monostearate, polysorbate 60, cetyl esters wax, cetearyl alcohol, 2-octyldodecanol, benzyl alcohol and water.

Topical application for the lower intestinal tract can be effected in a rectal suppository formulation (see below) or in a suitable enema formulation.

To be effective therapeutically as central nervous system targets, the compounds should readily penetrate the blood-brain barrier when peripherally administered. Compounds which cannot penetrate the blood-brain barrier can be effectively administered by an intraventricular route or other appropriate delivery system suitable for administration to the brain.

DOSAGE

The compounds and compositions of the present invention may be administered by a single dose, multiple discrete doses or continuous infusion. The compounds are well suited to continuous infusion. Pump means, particularly subcutaneous pump means, are preferred for continuous infusion.

Dose levels on the order of about 0.1 mg to about 10,000 mg of the active ingredient compound, but less than 40 mg/kg where the compound is Suramin, are useful in the treatment of the above conditions, with preferred levels being about 0.1 mg to about 1,000 mg. The specific dose level for any particular patient will vary depending upon a variety of factors, including the activity and the possible toxicity of the specific compound employed; the age, body weight, general health, sex and diet of the patient; the time of administration; the rate of excretion; drug combination; the severity of the particular disease being treated; and the form of administration. Typically, *in vitro* dosage-effect results provide useful guidance on the proper doses for patient administration. Studies in animal models are also helpful. The considerations for determining the proper dose levels are well known in the art.

The specific dose may be calculated according to considerations of body weight, body surface area or organ size. Further refinement of the calculations necessary to determine the appropriate dosage for treatment involving each of the above mentioned formulations is routinely made by those of ordinary skill in the art and is within the ambit of tasks routinely performed, especially in light of the dosage information and assays disclosed herein. Appropriate dosages may be determined using established assays in conjunction with appropriate dose-response data. One skilled in the art will appreciate that the dosage used in localized formulations of the invention normally will be smaller as compared to that used in a systemic injection or oral administration.

ADMINISTRATION REGIMEN

For the methods of the present invention, any

administration regimen well known to an ordinarily skilled artisan for regulating the timing and sequence of drug delivery can be used and repeated as necessary to effect treatment. Such regimen may include pretreatment and/or co-administration with additional therapeutic agents.

CO-ADMINISTRATION WITH OTHER TREATMENTS

The compounds and compositions of the present invention may be used alone or in combination with one or more additional agent(s) for simultaneous, separate or sequential use.

The additional agent(s) may be any therapeutic agent(s) known to an ordinarily skilled artisan, including without limitation: one or more compound(s) of the present invention; and one or more neurotrophic factor(s) selected from the group consisting of neurotrophic growth factor, brain derived growth factor, glial derived growth factor, ciliary neurotrophic factor, insulin growth factor, acidic fibroblast growth factor, basic fibroblast growth factor, platelet-derived growth factor, neurotrophin-3, neurotrophin-4 and neurotrophin-5; one or more neoplastic factors.

The compounds of the present invention can be co-administered with one or more therapeutic agents either (i) together in a single formulation, or (ii) separately in individual formulations designed for optimal release rates of their respective active agent. Each formulation may contain from about 0.01% to about 99.99% by weight, preferably from about 3.5% to about 60% by weight, of a compound of the present invention, as well as one or more pharmaceutical excipients, such as wetting, emulsifying and pH buffering agents.

EXAMPLES

The following examples are illustrative of the present invention and are not intended to be limitations thereon.

EXAMPLE 1

Synthesis of 4-Phenylbutyl 2-(3,3-dimethyl-2-oxopentanoyl)perhydropyridazinecarboxylate (Compound 15)

a. Synthesis of tert-butyl 2-[(tert-butyl)oxycarbonyl]-perhydropyridazinecarboxylate

A solution of di-Boc hydrazine (20 g, 84.4 mmol) in 150 DMF was added dropwise to a suspended solution of 6.75g (168.8 mmol) NaH in 75 ml DMF under nitrogen. After the mixture was stirred for 30 minutes at room temperature, a solution of 1,4 dibromobutane (18.2 g, 84.4 mmol) in 25 ml DMF was added dropwise. The reaction was allowed to stir overnight at room temperature. The reaction was then concentrated, followed by partition between 200 ml CH₂Cl₂ and 200 ml water. The aqueous layer was extracted with additional 200 ml CH₂Cl₂. The combined organic layers were dried over MgSO₄, and filtered and concentrated. The crude product was further purified by silica gel chromatography to yield 20.2 g (82% yield) product. The product was analyzed by GC/MS as pure compound with M⁺ 286.

b. Synthesis of perhydropyridazine

2.83 ml (36.7 mmol) TFA was added dropwise to a solution of tert-butyl 2-[(tert-butyl)oxycarbonyl]perhydropyridazinecarboxylate (1.5 g, 5.2 mmol) in 7 ml CH₂Cl₂, and the mixture was stirred overnight. At this time, the reaction was completed and 5.85 ml (42 mmol) triethylamine was added to quench the reaction. The reaction was concentrated and the residue, which contained product, was used without further

purification.

c. Synthesis of 4-phenylbutyl perhydropyridazine-carboxylate

A solution containing 1,1'-carbonyl diimidazole (0.893 g, 5.5 mmol) in 5 ml CH_2Cl_2 was added slowly to a solution of CH_2Cl_2 containing phenylbutyl alcohol (0.89 ml, 5.77 mmol). After stirring at room temperature for 1 hour, this solution was then added slowly to a solution containing perhydropyridazine mentioned above. The reaction was allowed to stir for overnight. The crude mixture was then concentrated and used without further purification.

d. Synthesis of methyl 2-oxo-2-{2-[(4-phenylbutyl)oxycarbonyl]perhydropyridazinyl} acetate

A solution of CH_2Cl_2 containing previous crude product of 4-phenylbutyl perhydropyridazinecarboxylate from last step was cooled to 0°C , and a solution of methyl oxalyl chloride (0.74 g, 5.77 mmol) in 5 ml CH_2Cl_2 was added dropwise over 0.5 hour. The resulting mixture was stirred at 0°C for 4 hours, and then warmed up to room temperature. The reaction mixture was diluted with 50 ml CH_2Cl_2 and washed with water. The organic layer was dried over MgSO_4 , and filtered and concentrated. The crude product was further purified by silica chromatography to yield 1.8 g (62% overall yield for three steps) product. ^1H NMR (CDCl_3 , 400 MHz): δ 1.39 (m, 2H); 1.69 (m, 6H); 2.62 (t, 2H, $J=8$); 2.83 (m, 1H); 3.10 (m, 1H); 3.79 (s, 3H); 4.16 (m, 3H); 4.31 (m, 1H); 7.22 (m, 5H).

e. Synthesis of 4-phenylbutyl 2-(3,3-dimethyl-2-oxopentanoyl)perhydropyridazine carboxylate

A solution of methyl 2-oxo-2-{2-[(4-phenylbutyl)-oxycarbonyl]perhydropyridazinyl} acetate (1.2 g, 3.45 mmol) in 15 ml dry THF was cooled to -78°C and treated with 5.2 ml of

1.0 M solution of 1,1-dimethylpropylmagnesium chloride in THF. After stirring the resulting homogeneous mixture at -78°C for 4 hours, the mixture was poured into saturated ammonium chloride (20 ml) and extracted into ethyl acetate. The organic layer was washed with water, dried and concentrated. The crude material was purified by silica gel column, eluting with 25% ethyl acetate in hexane, to obtain 0.98 g product (73% yield). $R_f = 0.73$ (2:1 hexane:EtOAc). ^1H NMR (CDCl_3 , 400 MHz): δ 0.81 (t, 3H, $J=7.1$); 1.13 (s, 3H); 1.20 (s, 3H); 1.64 (m, 10H); 2.64 (m, 2H); 2.86 (m, 1H); 3.20 (m, 1H); 3.99 (m, 1H); 4.19 (m, 2H); 4.35 (m, 1H); 7.24 (m, 5H). Anal. Calcd. for $\text{C}_{22}\text{H}_{32}\text{N}_2\text{O}_4$: C, 68.01; H, 8.30; N, 7.21. Found: C, 68.10; H, 8.29; N, 7.15.

EXAMPLE 2

Synthesis of 4-Phenylbutyl 2-[benzylsulfonyl]- perhydropyridazinecarboxylate (Compound 106)

A solution of α -toluene sulfonyl chloride (1.12 g, 5.77 mmol) in CH_2Cl_2 was added to a CH_2Cl_2 solution containing 4-phenylbutyl perhydropyridazinecarboxylate (1.37 g, 5.2 mmol) and triethylamine (0.83 ml, 6 mmol). The reaction was stirred overnight at room temperature under nitrogen atmosphere, and then diluted to 50 ml CH_2Cl_2 . The organic layer was washed with water, dried, and concentrated. The crude material was purified by silica gel column to yield 1.4 g (64%) final product as clear oil. $R_f = 0.60$ (2:1 hexane:EtOAc). ^1H NMR (CDCl_3 , 400 MHz): δ 1.68 (m, 8H); 2.67 (m, 2H); 2.90 (m, 1H); 3.38 (m, 2H); 4.22 (m, 5H); 7.32 (m, 10H). Anal. Calcd. for $\text{C}_{22}\text{H}_{28}\text{N}_2\text{S}_1\text{O}_4$: C, 63.44; H, 6.78; N, 6.73, S, 7.70. Found: C, 63.86; H, 6.83; N, 6.41, S, 7.58.

EXAMPLE 3

**Synthesis of 4-Phenylbutyl 2-(N-cyclohexylcarbamoyl)-
perhydropyridazinecarboxylate (Compound 153)**

Cyclohexylisocyanate (0.38 g, 3.0 mmol) was added to a CH₂Cl₂ solution containing 4-phenylbutyl perhydropyridazinecarboxylate (0.72 g, 2.75 mmol) and triethylamine (0.42 ml, 3 mmol). The reaction was stirred overnight at room temperature under nitrogen atmosphere, and then diluted to 50 ml CH₂Cl₂. The organic layer was washed with water, dried, and concentrated. The crude material was purified by silica gel column to yield 0.95 g (89%) final product as clear oil. R_f = 0.28 (2:1 hexane:EtOAc). ¹H NMR (CDCl₃, 400 MHz): δ 1.10 (m, 3H); 1.33 (m, 3H); 1.69 (m, 10H); 1.88 (m, 2H); 2.62 (m, 2H); 2.72 (m, 1H); 2.87 (m, 1H); 3.60 (m, 1H); 4.13 (m, 3H); 4.38 (m, 1H); 5.11 (d, 1H, J=8.3); 7.23 (m, 5H). Anal. Calcd. for C₂₂H₂₈N₃O₃ - 0.14H₂O: C, 67.75; H, 8.60; N, 10.77. Found: C, 67.75; H, 8.45; N, 10.90.

EXAMPLE 4

**Synthesis of 3,3-Dimethyl-1-[2-(6-phenylhexanoyl)perhydropyridazinyl]pentane-1,2-dione
(Compound 12) using Scheme 2**

a. Synthesis of (tert-butoxy)-N-[benzylamino]formamide

A solution of benzyl carbazate (25 g, 150.4 mmol), Boc anhydride (42.7 g, 195.5 mmol), triethylamine (19.8 g, 195.5 mmol), DMAP (0.9 g, 7.5 mmol) in 650 ml CH₂Cl₂ was stirred for 24 hours. The mixture was concentrated and purified by silica gel column, eluting with 20% ethyl acetate in hexane, to yield 36 g (90%) product. ¹H NMR (CDCl₃, 400 MHz): δ 1.40 (m, 9H); 5.25 (m, 2H); 7.36 (m, 5H).

b. Synthesis of tert-butyl 2-benzylperhydropyridazinecarboxylate

A solution of (tert-butoxy)-N-[benzylamino]formamide (35 g, 131 mmol) in 300 DMF was added dropwise to a suspended solution of 6.3 g (262 mmol) NaH in 130 ml DMF under nitrogen. After the mixture was stirred for 30 minutes at room temperature, a solution of 1,4 dibromobutane (28.4 g, 131 mmol) in 50 ml DMF was added dropwise. The reaction was allowed to stir overnight at room temperature. The reaction was then concentrated, followed by partition between 200 ml CH₂Cl₂ and 200 ml water. The aqueous layer was extracted with additional 200 ml CH₂Cl₂. The combined organic layers were dried over MgSO₄, and filtered and concentrated. The crude product was further purified by silica chromatography to yield 13.5 g (32% yield) product. ¹H NMR (CDCl₃, 400 MHz): δ 1.46 (m, 9H); 1.64 (m, 4H); 2.88 (m, 2H); 4.20 (m, 2H); 5.16 (m, 2H); 7.31 (m, 5H).

c. Synthesis of methyl 2-oxo-2-[2-benzylperhydropyridazinyl]acetate

20% TFA in CH₂Cl₂ was cooled to 0°C and added dropwise to a solution of tert-butyl 2-benzylperhydropyridazine-carboxylate (13.34 g, 41.7 mmol) in 10 ml CH₂Cl₂. The mixture was stirred overnight. At this time, the mixture was cooled to 0°C and 12.66 ml (125 mmol) triethylamine was added, followed by addition dropwise of methyl oxalyl chloride (5.62, 45.9 mmol) in 5 ml CH₂Cl₂. The mixture was allowed to stirred 2 hours at 0°C and warmed up to room temperature overnight. The reaction was diluted with addition of CH₂Cl₂ and washed with water. The organic layer was dried over MgSO₄, and filtered and concentrated. The crude product was further purified by silica chromatography to yield 9.2 g (72.4% yield) product as clear oil. ¹H NMR (CDCl₃, 400 MHz): δ 1.72 (m, 4H); 2.85 (m, 1H); 3.12 (m, 1H); 3.67 (s, 3H); 4.15 (m, 1H);

4.35 (m, 1H); 5.20 (m, 2H); 7.35 (m, 5H).

d. Synthesis of 3,3-dimethyl-1-[2-benzylperhydro-pyridazinyl]pentane-1,2-dione

A solution of methyl 2-oxo-2-[2-benzylperhydro-pyridazinyl]acetate (9.0 g, 29.4 mmol) in 30 ml dry THF was cooled to -78°C and treated with 35 ml of 1.0 M solution of 1,1-dimethylpropylmagnesium chloride in THF. After stirring the resulting homogeneous mixture at for 5 hours, the mixture was poured into saturated ammonium chloride (150 ml) and extracted into ethyl acetate. The organic layer was washed with water, dried and concentrated. The crude material was purified by silica gel column, eluting with 10% ethyl acetate in hexane, to obtain 7.0 g product (69% yield) as clear oil. ¹H NMR (CDCl₃, 400 MHz): δ 0.76 (t, 3H, J=7.0); 1.06 (s, 6H); 1.69 (m, 6H); 2.80 (m, 1H); 3.15 (m, 1H), 4.03 (m, 1H); 4.13 (m, 1H), 5.18 (m, 2H), 7.36 (m, 5H).

e. Synthesis of 3,3-dimethyl-1-perhydropyridazinyl-pentane-1,2-dione

1 g 10% Pd/C was added to a solution of 3,3-dimethyl-1-[2-benzylperhydropyridazinyl] pentane-1,2-dione (7.0 g, 20.2 mmol) in 70 ml EtOH. The mixture was under hydrogenation at room pressure (1 atm) overnight. The product was obtained as white solid after filtering Pd catalyst and concentration (3.8 g, 89%). ¹H NMR (CDCl₃, 400 MHz): δ 0.88 (t, 3H, J=7.0); 1.19 (s, 6H); 1.65 (m, 4H); 1.79 (m, 2H); 2.85 (m, 2H); 3.42 (m, 1H); 3.56 (m, 1H).

f. Synthesis of 3,3-dimethyl-1-[2-(6-phenylhexanoyl)-perhydropyridazinyl]pentane-1,2-dione

To a solution of 5-phenylvaleric acid (0.2 g, 1.1 mmol) in 3 ml CH₂Cl₂ was added triethylamine (0.15 ml, 1.1 mmol),

followed by isobutyl chloroformate (0.15 g, 1.1 mmol) at 0°C. After stirring for 5 minutes, a solution of 3,3-dimethyl-1-perhydropyridazinylpentane-1,2-dione (0.212 g, 1 mmol) in 1 ml CH₂Cl₂ was added. The reaction was gradually warmed up to room temperature. The crude material was subject to silica gel purification to yield final product as clear oil (0.20 g, 55%). R_f = 0.58 (33% EtOAc/hexane). ¹H NMR (CDCl₃, 400 MHz): δ 0.89 (t, 3H, J=7.5); 1.24 (s, 6H); 1.37 (m, 2H); 1.68 (m, 6H); 1.74 (m, 4H); 2.23 (m, 2H); 2.62 (t, 2H, J=7.60); 2.80 (m, 2H); 4.53 (m, 2H); 7.21 (m, 5H). Anal. Calcd. for C₂₃H₃₄N₂O₃: C, 71.47; H, 8.87; N, 7.25. Found: C, 71.54; H, 8.80; N, 7.32.

EXAMPLE 5

Synthesis of 3,3-Dimethyl-1-[2-(6-(3-pyridyl)hexanoyl)-perhydropyridazinyl]-pentane-1,2-dione (Compound 13) using

Scheme 2

a. Synthesis of 1-(2-hex-5-ynoylperhydropyridazinyl)-3,3-dimethylpentane-1,2-dione

To a solution of 5-hexynoic acid (0.467 g, 4 mmol) in 10 ml CH₂Cl₂ was added triethylamine (0.56 ml, 4 mmol), followed by isobutyl chloroformate (0.53 ml, 4 mmol) at 0°C. After stirring for 5 minutes, a solution of 3,3-dimethyl-1-perhydropyridazinyl pentane-1,2-dione (0.424 g, 2 mmol) in 1 ml CH₂Cl₂ was added. The reaction was gradually warmed up to room temperature. The crude material was subject to silica gel purification to yield final product as clear oil (0.385 g, 63%). ¹H NMR (CDCl₃, 400 MHz): δ 0.91 (t, 3H, J=7.0); 1.26 (s, 6H); 1.76 (m, 8H); 2.28 (m, 2H); 2.50 (m, 2H); 2.88 (m, 2H); 3.60 (m, 1H); 4.50 (m, 2H).

b. Synthesis of 3,3-dimethyl-1-[2-(6-(3-pyridyl)hex-5-

ynoyl)perhydropyridazinyl] pentane-1,2-dione

To a solution of 1-(2-hex-5-ynoylperhydropyridazinyl)-3,3-dimethylpentane-1,2-dione (0.384 g, 1.25 mmol) in 10 ml CH₂Cl₂ under nitrogen was added 3-iodopyridine (0.283 g, 1.38 mmol), (Ph₃P)₂PdCl₂ (0.044 g, 0.06 mmol), CuI (0.0024 g, 0.013 mmol) and triethylamine (0.3 ml, 2 mmol). The reaction mixture was stirred 30 minutes at room temperature and then refluxed overnight. The mixture was concentrated and purified by silica gel column, eluting with 30% ethyl acetate in hexane, to yield product as light yellow oil (0.31 g, 65%).
¹H NMR (CDCl₃, 400 MHz): δ 0.84 (t, 3H, J=7.4); 1.21 (s, 6H); 1.70 (m, 6H); 1.96 (m, 2H), 2.52 (m, 3H); 2.90 (m, 2H); 3.60 (m, 1H); 4.42 (m, 2H); 7.20 (m, 1H); 7.66 (m, 1H); 8.49 (m, 1H); 8.62 (m, 1H).

c. Synthesis of 3,3-dimethyl-1-[2-(6-(3-pyridyl)-hexanoyl)perhydropyridazinyl] pentane-1,2-dione

0.1 g PtO₂ was added to a solution of 3,3-dimethyl-1-[2-(6-(3-pyridyl)hex-5 ynoyl)perhydropyridazinyl] pentane-1,2-dione (0.3 g, 0.8 mmol) in 20 ml dry MeOH. The mixture was under hydrogenation at room pressure (1 atm) overnight. The product was obtained as clear oil after filtering the catalyst, concentration and purifying on a silica gel (0.125 g, 41%). R_f = 0.18 (EtOAc). ¹H NMR (CDCl₃, 400 MHz): δ 0.89 (t, 3H, J=7.4); 1.24 (s, 6H); 1.38 (m, 2H); 1.66 (m, 10H); 2.14 (m, 2H); 2.63 (m, 2H); 2.82 (m, 2H); 4.60 (m, 2H); 7.23 (m, 4H). Anal. Calcd. for C₂₂H₃₃N₃O₃: C, 68.19; H, 8.58; N, 10.84. Found: C, 68.40; H, 8.52; N, 10.62.

EXAMPLE 6

Synthesis of 2-Cyclohexyl-2,5,6,7,8,8a-hexahydro-2,8a-diazaindolizine-1,3-dione using Scheme 1

To a solution of 4-phenylbutyl 2-(N-cyclohexyl-carbamoyl)perhydropyridazine carboxylate (0.53 g, 1.37 mmol) in 5 ml THF at 0°C under nitrogen was added 1.37 ml of 1 M LHMDs in THF. The mixture was allowed to stir overnight, gradually warming up to room temperature. The mixture was concentrated and purified by silica gel column, eluting with 30% ethyl acetate in hexane, to afford product (0.27 g, 83%). $R_f = 0.32$ (2:1 hexane:EtOAc). ^1H NMR (CDCl_3 , 400 MHz): δ 1.27(m, 3H); 1.75(m, 9H); 2.12(m, 2H); 3.50(m, 4H); 3.87(m, 1H). Anal. Calcd. for $\text{C}_{12}\text{H}_{19}\text{N}_3\text{O}_2$: C, 60.74; H, 8.07; N, 17.71. Found: C, 60.61; H, 8.11; N, 17.82.

EXAMPLE 7

Synthesis of Compounds 1, 2, 5, 6 and 18

Compounds 1, 2, 5, 6 and 18 were synthesized by the general method illustrated in Scheme 2 and exemplified in Example 4.

1) 3,3-Dimethyl-N-[2-(5-phenylpentanoyl)tetrahydro-1H-1-pyrazolyl]-1,2-pentane-dione. $R_f = 0.25$ (2:1 hexane:EtOAc). ^1H NMR (CDCl_3 , 300 MHz): δ 0.81-0.83 (m, 3H); 1.14 (s, 6H); 1.21 (m, 2H); 1.55-1.62 (m, 8H); 2.02 (m, 2H); 2.61 (m, 4H); 7.14-7.28 (m, 5H). Anal. Calcd. for $\text{C}_{21}\text{H}_{30}\text{N}_2\text{O}_3$: C, 70.36; H, 8.44; N, 7.81. Found: C, 70.10; H, 8.41; N, 7.77.

2) 3,3-Dimethyl-N-[2-(3-phenylpropanoyl)tetrahydro-1H-1-pyrazolyl]-1,2-pentanedione. $R_f = 0.60$ (2:1 hexane:EtOAc). ^1H NMR (CDCl_3 , 300 MHz): δ 0.80-0.85 (t, 3H); 1.11-1.15 (m, 8H); 1.58-2.02 (m, 6H); 2.50-2.95 (m, 4H); 7.17-7.28 (m, 5H). Anal. Calcd. for $\text{C}_{19}\text{H}_{26}\text{N}_2\text{O}_3$: C, 69.06; H, 7.93; N, 8.48. Found: C, 68.98; H, 7.90; N, 8.41.

5) 3,3-Dimethyl-1-[2-(4-phenylbutanoyl)pyrazolidinyl]-

pentane-1,2-dione. $R_f = 0.5$ (Hexane:EtAc 1:1). ^1H NMR (CDCl_3 , 400 MHz): δ 0.87 (t, 3H, $J=7.5$); 1.22 (s, 3H); 1.26 (s, 3H); 1.64 (m, 2H); 1.92-2.07 (m, 5H), 2.20 (m, 1H), 2.63 (m, 2H); 3.25 (m, 2H); 3.80 (m, 2H); 7.27 (m, 5H, aromatic). Anal. Calcd. for $\text{C}_{20}\text{H}_{28}\text{N}_2\text{O}_3$: C, 69.05 (69.02); H, 8.27 (8.22); N, 8.06 (8.05).

6) 3,3-Dimethyl-1-[2-(6-phenylhexanoyl)pyrazolidinyl]-pentane-1,2-dione. $R_f = 0.5$ (Hexane:EtAc 1:1). ^1H NMR (CDCl_3 , 400 MHz): δ 0.87 (t, 3H, $J=7.5$); 1.22 (s, 3H); 1.26 (s, 3H); 1.35 (m, 2H); 1.59 (m, 6H); 2.07 (m, 2H), 2.20 (m, 1H), 2.60 (m, 3H); 3.25 (m, 2H); 3.70 (m, 2H); 7.26 (m, 5H, aromatic). Anal. Calcd. for $\text{C}_{24}\text{H}_{32}\text{N}_2\text{O}_3$: C, 70.65 (70.94); H, 8.70 (8.66); N, 7.36 (7.52).

18) 3,3-Dimethyl-1-[2-(5-phenyl)pentanoyl]perhydropyridazinyl]pentane-1,2-dione. $R_f = 0.53$ (33% EtOAc/hexane). ^1H NMR (CDCl_3 , 400 MHz): δ 0.90 (t, 3H, $J=7.44$); 1.24 (s, 6H); 1.64 (m, 8H); 2.28 (m, 2H); 2.65 (m, 2H); 2.80 (m, 2H); 3.12 (m, 1H); 3.58 (m, 1H); 4.54 (m, 2H); 7.22 (m, 5H). Anal. Calcd. for $\text{C}_{22}\text{H}_{32}\text{N}_2\text{O}_3$: C, 70.94; H, 8.66; N, 7.52. Found: C, 71.07; H, 8.59; N, 7.51.

EXAMPLE 8

Synthesis of Compounds 3, 4 and 7

Compounds 3, 4 and 7 were synthesized by the general method illustrated in Scheme 2 and exemplified by Example 5.

3) 3,3-Dimethyl-1-[2-(5-(3-pyridyl)pent-4-ynoyl)-pyrazolidinyl]pentane-1,2-dione. $R_f = 0.2$ (EtOAc). ^1H NMR (CDCl_3 , 400 MHz): δ 0.87 (t, 3H, $J=7.5$); 1.22 (s, 3H); 1.26 (s, 3H); 1.63 (m, 2H); 2.1 (m, 2H); 2.73 (m, 4H); 3.20-3.85 (m, 4H); 7.19 (m, 1H); 7.66 (m, 1H); 8.5 (m, 2H). Anal. Calcd. for $\text{C}_{20}\text{H}_{25}\text{N}_3\text{O}_3$: C, 67.67 (64.58), H: 6.91 (7.09), N: 10.63

(10.82).

4) 3,3-Dimethyl-1-[2-(5-(cyano)pent-4-ynoyl)pyrazolidinyl]-
pentane-1,2-dione. $R_f = 0.45$ (EtAc). ^1H NMR (CDCl_3 , 400 MHz):
 δ 0.87 (t, $J=7.5$); 0.90 (m, 2H); 1.22 (s, 3H); 1.26 (s, 3H);
 1.64 (m, 2H); 2.03-2.20 (m, 3H), 2.52 (m, 2H), 2.63 (m, 1H);
 3.69 (m, 3H). Anal. Calcd. for $\text{C}_{15}\text{H}_{22}\text{N}_2\text{O}_3$: C, 64.55 (64.73);
 H, 7.98 (7.97); N, 9.98 (10.06).

7) 3,3-Dimethyl-1-[2-(5-(3-pyridyl)pentanoyl)-
pyrazolidinyl]pentane-1,2-dione. $R_f = 0.3$ (EtAc). ^1H NMR
 (CDCl_3 , 400 MHz): δ 0.87 (t, 3H, $J=7.5$); 1.22 (s, 3H); 1.26
 (s, 3H); 1.37 (m, 2H); 1.65 (m, 6H); 2.1 (m, 2H); 2.30 (m,
 1H); 2.62 (m, 3H); 3.20-3.85 (m, 4H); 7.19 (m, 1H); 7.66 (m,
 1H); 8.5 (m, 2H). Anal. Calcd. for $\text{C}_{20}\text{H}_{29}\text{N}_3\text{O}_3$: C, 65.74
 (65.98); H, 8.06 (8.20); N, 11.09 (10.89).

EXAMPLE 9

Synthesis of Compounds 8-10, 11, 14, 16 and 17

Compounds 8-10, 13, 16, 18 and 21 were synthesized by the
 general method illustrated in Scheme 1 and exemplified in
 Example 1.

8) 3-Phenylpropyl 2-(3,3-dimethyl-2-oxopentanoyl)-
pyrazolidinecarboxylate. $R_f = 0.4$ (25% EtOAc/Hexane). ^1H NMR
 (CDCl_3 , 400 MHz): δ 0.86 (t, 3H, $J=7.4$); 1.22 (s, 6H); 1.66
 (t, 2H, $J=7.5$); 2.00-2.12 (m, 4H); 2.72 (t, 2H, $J=7.4$); 3.60
 (br s, 4H); 4.18 (t, 2H, $J=6.5$); 7.18-7.31 (m, 5H). Anal.
 Calcd. for $\text{C}_{20}\text{H}_{28}\text{N}_2\text{O}_4$: C, 66.64; H, 7.83; N, 7.77. Found: C,
 66.73; H, 7.81; N, 7.72.

9) 3-(3-Pyridyl)propyl 2-(3,3-dimethyl-2-oxopentanoyl)-
pyrazolidinecarboxylate. $R_f = 0.1$ (100% EtOAc). ^1H NMR
 (CDCl_3 , 400 MHz): δ 0.85 (t, 3H, $J=7.5$); 1.21 (s, 6H); 1.67

(t, 2H, J=7.5); 2.00-2.13 (m, 4H); 2.72 (t, 2H, J=7.5); 3.62 (br s, 4H); 4.19 (t, 2H, J=6.4); 7.28 (br s, 1H), 7.54 (d, 1H, J=7.7); 8.48 (s, 2H). Anal. Calcd. for $C_{19}H_{27}N_3O_4 - 0.35 H_2O$: C, 62.06; H, 7.59; N, 11.43. Found: C, 61.77; H, 7.53; N, 11.36.

10) 4-Phenylbutyl 2-(3,3-dimethyl-2-oxopentanoyl)-pyrazolidinecarboxylate. $R_f = 0.6$ (25% EtOAc/Hexane). 1H NMR ($CDCl_3$, 400 MHz): δ 0.83 (t, 3H, J=7.5); 1.19 (s, 6H); 1.67 (t, 2H, J=7.5); 1.60-1.69 (m, 4H); 2.07 (t, 2H, J=7.4); 2.62 (t, 2H, J=6.4); 3.60 (br s, 4H); 4.13 (t, 2H, J=6.1); 7.28-7.15 (m, 5H). Anal. Calcd. for $C_{21}H_{30}N_2O_4$: C, 67.35; H, 8.07; N, 7.48. Found: C, 67.54; H, 8.31; N, 7.40.

11) 2-Phenylethyl 2-(3,3-dimethyl-2-oxopentanoyl)-pyrazolidinecarboxylate. $R_f = 0.5$ (25% EtOAc/Hexane). 1H NMR ($CDCl_3$, 400 MHz): δ 0.83 (t, 3H, J=7.5); 1.18 (s, 6H); 1.63 (m, 2H); 1.99 (m, 2H); 2.97 (t, 2H, J=7.1); 3.60 (br s, 4H); 4.35 (t, 2H, J=6.6); 7.19-7.30 (m, 5H). Anal. Calcd. for $C_{17}H_{26}N_2O_4$: C, 65.88; H, 7.56; N, 8.09. Found: C, 65.82; H, 7.51; N, 8.02.

14) 3-phenylpropyl 2-(3,3-dimethyl-2-oxopentanoyl)-perhydropyridazinecarboxylate. $R_f = 0.73$ (2:1 hexane:EtOAc). 1H NMR ($CDCl_3$, 400 MHz): 0.82 (t, 3H, J=7.4); 1.16 (s, 3H); 1.22 (s, 3H); 1.67 (m, 6H); 2.00 (m, 2H); 2.69 (t, 2H, J=7.9); 2.86 (m, 1H); 3.23 (m, 1H); 4.00 (m, 1H); 4.20 (m, 2H); 4.37 (m, 1H); 7.23 (m, 5H). Anal. Calcd. for $C_{21}H_{30}N_2O_4$: C, 67.35; H, 8.07; N, 7.48. Found: C, 67.51; H, 8.11; N, 7.39.

16) 5-Phenylpentyl 2-(3,3-dimethyl-2-oxopentanoyl)-perhydropyridazinecarboxylate. $R_f = 0.74$ (2:1 hexane:EtOAc). 1H NMR ($CDCl_3$, 400 MHz): δ 0.82 (t, 3H, J=7.4); 1.14 (s, 3H); 1.21 (s, 3H); 1.38 (m, 2H); 1.65 (m, 10H); 2.62 (t, 2H, J=7.6); 2.83 (m, 1H); 3.20 (m, 1H); 3.98 (m, 1H); 4.15 (m,

2H); 4.33 (m, 1H); 7.23 (m, 5H). Anal. Calcd. for $C_{23}H_{34}N_2O_4$: C, 68.63; H, 8.51; N, 6.96. Found: C, 68.70; H, 8.47; N, 7.08.

17) 4-(3-Pyridyl)butyl 2-(3,3-dimethyl-2-oxopentanoyl)-
 5 perhydropyridazinecarboxylate. $R_f = 0.45$ (100% EtOAc). 1H NMR (CDCl₃, 400 MHz): δ 0.81 (t, 3H, J=7.5); 1.14 (s, 3H); 1.20 (s, 3H); 1.70 (m, 10H); 2.66 (m, 2H); 2.86 (m, 1H); 3.20 (m, 1H); 4.00 (m, 1H); 4.18 (m, 2H); 4.36 (m, 1H); 7.22 (m, 1H); 7.50 (m, 1H); 8.45 (m, 2H). Anal. Calcd. for $C_{21}H_{31}N_3O_4$ - 0.14
 10 H₂O: C, 64.34; H, 8.04; N, 10.72. Found: C, 64.34; H, 8.02; N, 10.83.

EXAMPLE 10

Synthesis of Compound 105

Compound 105 was synthesized by the general method
 15 illustrated in Scheme 1 and exemplified in Example 2.

3-Phenylpropyl 2-[benzylsulfonyl]pyrazolidine-carboxylate. $R_f = 0.5$ (40% EtOAc/Hexane). 1H NMR (CDCl₃, 400 MHz): δ 2.01-2.17 (m, 4H); 2.72 (t, 2H, J=7.8); 3.68 (br s, 4H); 4.23 (t, 2H, J=6.6); 4.51 (s, 2H); 7.17-7.50 (m, 10H). Anal. Calcd.
 20 for $C_{20}H_{24}N_2SO_4$: C, 61.83; H, 6.23; N, 7.21; S, 8.25. Found: C, 61.63; H, 6.21; N, 7.05; S, 8.07.

EXAMPLE 11

Synthesis of Compound 153

Compound 153 was synthesized by the general method
 25 illustrated in Scheme 1 and exemplified in Example 3.

3-Phenylpropyl 2-(N-cyclohexylcarbamoyl)pyrazolidine-carboxylate. $R_f = 0.5$ (60% EtOAc/Hexane). 1H NMR (CDCl₃, 400 MHz): δ 1.09-2.00 (m, 15H); 2.69 (t, 2H, J=7.8); 3.70 (br s, 4H); 4.18 (t, 2H, J=6.4); 5.46 (d, 1H, J=8.2); 7.16-7.30 (m,

5H). Anal. Calcd. for $C_{20}H_{29}N_3O_3$: C, 66.83; H, 8.13; N, 11.69. Found: C, 66.73; H, 8.28; N, 11.59.

EXAMPLE 12

K_i Test

5 Inhibition of the peptidyl-prolyl isomerase (rotamase) activity of the inventive compounds can be evaluated by known methods described in the literature (Harding et al., *Nature*, 1989, 341:758-760; Holt et al. *J. Am. Chem. Soc.*, 115:9923-9938). These values are obtained as apparent K_i 's and are
10 presented for representative compounds in Table V. The *cis-trans* isomerization of an alanine-proline bond in a model substrate, N-succinyl-Ala-Ala-Pro-Phe-*p*-nitroanilide, is monitored spectrophotometrically in a chymotrypsin-coupled assay, which releases *para*-nitroanilide from the *trans* form of
15 the substrate. The inhibition of this reaction caused by the addition of different concentrations of inhibitor is determined, and the data is analyzed as a change in first-order rate constant as a function of inhibitor concentration to yield the apparent K_i values.

20 In a plastic cuvette are added 950 μ L of ice cold assay buffer (25 mM HEPES, pH 7.8, 100 mM NaCl), 10 μ L of FKBP (2.5 mM in 10 mM Tris-Cl pH 7.5, 100 mM NaCl, 1 mM dithiothreitol), 25 μ L of chymotrypsin (50 mg/ml in 1 mM HCl) and 10 μ L of test compound at various concentrations in dimethyl sulfoxide. The
25 reaction is initiated by the addition of 5 μ L of substrate (succinyl-Ala-Phe-Pro-Phe-*para*-nitroanilide, 5 mg/mL in 2.35 mM LiCl in trifluoroethanol).

30 The absorbance at 390 nm versus time is monitored for 90 seconds using a spectrophotometer and the rate constants are determined from the absorbance versus time data files.

The results of these experiments are presented in TABLE V under the column "Ki".





EXAMPLE 13





MPTP Model of Parkinson's Disease





5 The neurotrophic and neuroregenerative effects of the
inventive compounds were demonstrated in an animal model of
neurodegenerative disease. MPTP lesioning of dopaminergic
neurons in mice was used as an animal model of Parkinson's
Disease. Four week old male CD1 white mice were dosed i.p.
10 with 30 mg/kg of MPTP for 5 days. Test compounds (4 or 10
mg/kg) or vehicle, were administered s.c. along with the MPTP
for 5 days, as well as for an additional 5 days following
cessation of MPTP treatment. At 18 days following MPTP
treatment, the animals were sacrificed and the striata were
15 dissected and homogenized. Immunostaining was performed on
sagittal and coronal brain sections using anti-tyrosine
hydroxylase 1 g to quantitate survival and recovery of
dopaminergic neurons. In animals treated with MPTP and
vehicle, a substantial loss of functional dopaminergic
20 terminals was observed as compared to non-lesioned animals.
Lesioned animals receiving test compounds showed a significant
recovery of TH-stained dopaminergic neurons.




The results of these experiments are presented in TABLE V under the column "% TH recovery".

TABLE V

No.	Structure	Ki nM	% TH recov 4 mg/kg s.c.	% TH recov 10 mg/kg p.o.
1		1175	14	
2				10
3				26
5				4

No.	Structure	Ki nM	% TH recov 4 mg/kg s.c.	% TH recov 10 mg/kg p.o.
6				32
7				57
12				35
13		3208		18

No.	Structure	Ki nM	% TH recov 4 mg/kg s.c.	% TH recov 10 mg/kg p.o.
14				3
15				21
16				49
17				38

No.	Structure	Ki nM	% TH recov 4 mg/kg s.c.	% TH recov 10 mg/kg p.o.
18				18
106				23
154				46

EXAMPLE 14

A patient is suffering from a disease, disorder or condition described above. The patient may then be administered an effective amount of a compound of the present invention. It is expected that after such treatment, the patient would not suffer any significant injury due to, would be protected from further injury due to, or would recover from the disease, disorder or condition.

The invention being thus described, it will be obvious that the same may be varied in many ways. Such variations are not to be regarded as a departure from the spirit and scope of the invention and all such modifications are intended to be included within the scope of the following claims.

5

100 101 102 103 104 105 106 107 108 109 110 111 112 113 114 115 116 117 118 119 120 121 122 123 124 125 126 127 128 129 130 131 132 133 134 135 136 137 138 139 140 141 142 143 144 145 146 147 148 149 150 151 152 153 154 155 156 157 158 159 160 161 162 163 164 165 166 167 168 169 170 171 172 173 174 175 176 177 178 179 180 181 182 183 184 185 186 187 188 189 190 191 192 193 194 195 196 197 198 199 200 201 202 203 204 205 206 207 208 209 210 211 212 213 214 215 216 217 218 219 220 221 222 223 224 225 226 227 228 229 230 231 232 233 234 235 236 237 238 239 240 241 242 243 244 245 246 247 248 249 250 251 252 253 254 255 256 257 258 259 260 261 262 263 264 265 266 267 268 269 270 271 272 273 274 275 276 277 278 279 280 281 282 283 284 285 286 287 288 289 290 291 292 293 294 295 296 297 298 299 300 301 302 303 304 305 306 307 308 309 310 311 312 313 314 315 316 317 318 319 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 409 410 411 412 413 414 415 416 417 418 419 420 421 422 423 424 425 426 427 428 429 430 431 432 433 434 435 436 437 438 439 440 441 442 443 444 445 446 447 448 449 450 451 452 453 454 455 456 457 458 459 460 461 462 463 464 465 466 467 468 469 470 471 472 473 474 475 476 477 478 479 480 481 482 483 484 485 486 487 488 489 490 491 492 493 494 495 496 497 498 499 500 501 502 503 504 505 506 507 508 509 510 511 512 513 514 515 516 517 518 519 520 521 522 523 524 525 526 527 528 529 530 531 532 533 534 535 536 537 538 539 540 541 542 543 544 545 546 547 548 549 550 551 552 553 554 555 556 557 558 559 560 561 562 563 564 565 566 567 568 569 570 571 572 573 574 575 576 577 578 579 580 581 582 583 584 585 586 587 588 589 590 591 592 593 594 595 596 597 598 599 600 601 602 603 604 605 606 607 608 609 610 611 612 613 614 615 616 617 618 619 620 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 648 649 650 651 652 653 654 655 656 657 658 659 660 661 662 663 664 665 666 667 668 669 670 671 672 673 674 675 676 677 678 679 680 681 682 683 684 685 686 687 688 689 690 691 692 693 694 695 696 697 698 699 700 701 702 703 704 705 706 707 708 709 710 711 712 713 714 715 716 717 718 719 720 721 722 723 724 725 726 727 728 729 730 731 732 733 734 735 736 737 738 739 740 741 742 743 744 745 746 747 748 749 750 751 752 753 754 755 756 757 758 759 760 761 762 763 764 765 766 767 768 769 770 771 772 773 774 775 776 777 778 779 780 781 782 783 784 785 786 787 788 789 790 791 792 793 794 795 796 797 798 799 800 801 802 803 804 805 806 807 808 809 810 811 812 813 814 815 816 817 818 819 820 821 822 823 824 825 826 827 828 829 830 831 832 833 834 835 836 837 838 839 840 841 842 843 844 845 846 847 848 849 850 851 852 853 854 855 856 857 858 859 860 861 862 863 864 865 866 867 868 869 870 871 872 873 874 875 876 877 878 879 880 881 882 883 884 885 886 887 888 889 890 891 892 893 894 895 896 897 898 899 900 901 902 903 904 905 906 907 908 909 910 911 912 913 914 915 916 917 918 919 920 921 922 923 924 925 926 927 928 929 930 931 932 933 934 935 936 937 938 939 940 941 942 943 944 945 946 947 948 949 950 951 952 953 954 955 956 957 958 959 960 961 962 963 964 965 966 967 968 969 970 971 972 973 974 975 976 977 978 979 980 981 982 983 984 985 986 987 988 989 990 991 992 993 994 995 996 997 998 999 1000 1001 1002 1003 1004 1005 1006 1007 1008 1009 1010 1011 1012 1013 1014 1015 1016 1017 1018 1019 1020 1021 1022 1023 1024 1025 1026 1027 1028 1029 1030 1031 1032 1033 1034 1035 1036 1037 1038 1039 1040 1041 1042 1043 1044 1045 1046 1047 1048 1049 1050 1051 1052 1053 1054 1055 1056 1057 1058 1059 1060 1061 1062 1063 1064 1065 1066 1067 1068 1069 1070 1071 1072 1073 1074 1075 1076 1077 1078 1079 1080 1081 1082 1083 1084 1085 1086 1087 1088 1089 1090 1091 1092 1093 1094 1095 1096 1097 1098 1099 1100 1101 1102 1103 1104 1105 1106 1107 1108 1109 1110 1111 1112 1113 1114 1115 1116 1117 1118 1119 1120 1121 1122 1123 1124 1125 1126 1127 1128 1129 1130 1131 1132 1133 1134 1135 1136 1137 1138 1139 1140 1141 1142 1143 1144 1145 1146 1147 1148 1149 1150 1151 1152 1153 1154 1155 1156 1157 1158 1159 1160 1161 1162 1163 1164 1165 1166 1167 1168 1169 1170 1171 1172 1173 1174 1175 1176 1177 1178 1179 1180 1181 1182 1183 1184 1185 1186 1187 1188 1189 1190 1191 1192 1193 1194 1195 1196 1197 1198 1199 1200 1201 1202 1203 1204 1205 1206 1207 1208 1209 1210 1211 1212 1213 1214 1215 1216 1217 1218 1219 1220 1221 1222 1223 1224 1225 1226 1227 1228 1229 1230 1231 1232 1233 1234 1235 1236 1237 1238 1239 1240 1241 1242 1243 1244 1245 1246 1247 1248 1249 1250 1251 1252 1253 1254 1255 1256 1257 1258 1259 1260 1261 1262 1263 1264 1265 1266 1267 1268 1269 1270 1271 1272 1273 1274 1275 1276 1277 1278 1279 1280 1281 1282 1283 1284 1285 1286 1287 1288 1289 1290 1291 1292 1293 1294 1295 1296 1297 1298 1299 1300 1301 1302 1303 1304 1305 1306 1307 1308 1309 1310 1311 1312 1313 1314 1315 1316 1317 1318 1319 1320 1321 1322 1323 1324 1325 1326 1327 1328 1329 1330 1331 1332 1333 1334 1335 1336 1337 1338 1339 1340 1341 1342 1343 1344 1345 1346 1347 1348 1349 1350 1351 1352 1353 1354 1355 1356 1357 1358 1359 1360 1361 1362 1363 1364 1365 1366 1367 1368 1369 1370 1371 1372 1373 1374 1375 1376 1377 1378 1379 1380 1381 1382 1383 1384 1385 1386 1387 1388 1389 1390 1391 1392 1393 1394 1395 1396 1397 1398 1399 1400 1401 1402 1403 1404 1405 1406 1407 1408 1409 1410 1411 1412 1413 1414 1415 1416 1417 1418 1419 1420 1421 1422 1423 1424 1425 1426 1427 1428 1429 1430 1431 1432 1433 1434 1435 1436 1437 1438 1439 1440 1441 1442 1443 1444 1445 1446 1447 1448 1449 1450 1451 1452 1453 1454 1455 1456 1457 1458 1459 1460 1461 1462 1463 1464 1465 1466 1467 1468 1469 1470 1471 1472 1473 1474 1475 1476 1477 1478 1479 1480 1481 1482 1483 1484 1485 1486 1487 1488 1489 1490 1491 1492 1493 1494 1495 1496 1497 1498 1499 1500 1501 1502 1503 1504 1505 1506 1507 1508 1509 1510 1511 1512 1513 1514 1515 1516 1517 1518 1519 1520 1521 1522 1523 1524 1525 1526 1527 1528 1529 1530 1531 1532 1533 1534 1535 1536 1537 1538 1539 1540 1541 1542 1543 1544 1545 1546 1547 1548 1549 1550 1551 1552 1553 1554 1555 1556 1557 1558 1559 1560 1561 1562 1563 1564 1565 1566 1567 1568 1569 1570 1571 1572 1573 1574 1575 1576 1577 1578 1579 1580 1581 1582 1583 1584 1585 1586 1587 1588 1589 1590 1591 1592 1593 1594 1595 1596 1597 1598 1599 1600 1601 1602 1603 1604 1605 1606 1607 1608 1609 1610 1611 1612 1613 1614 1615 1616 1617 1618 1619 1620 1621 1622 1623 1624 1625 1626 1627 1628 1629 1630 1631 1632 1633 1634 1635 1636 1637 1638 1639 1640 1641 1642 1643 1644 1645 1646 1647 1648 1649 1650 1651 1652 1653 1654 1655 1656 1657 1658 1659 1660 1661 1662 1663 1664 1665 1666 1667 1668 1669 1670 1671 1672 1673 1674 1675 1676 1677 1678 1679 1680 1681 1682 1683 1684 1685 1686 1687 1688 1689 1690 1691 1692 1693 1694 1695 1696 1697 1698 1699 1700 1701 1702 1703 1704 1705 1706 1707 1708 1709 1710 1711 1712 1713 1714 1715 1716 1717 1718 1719 1720 1721 1722 1723 1724 1725 1726 1727 1728 1729 1730 1731 1732 1733 1734 1735 1736 1737 1738 1739 1740 1741 1742 1743 1744 1745 1746 1747 1748 1749 1750 1751 1752 1753 1754 1755 1756 1757 1758 1759 1760 1761 1762 1763 1764 1765 1766 1767 1768 1769 1770 1771 1772 1773 1774 1775 1776 1777 1778 1779 1780 1781 1782 1783 1784 1785 1786 1787 1788 1789 1790 1791 1792 1793 1794 1795 1796 1797 1798 1799 1800 1801 1802 1803 1804 1805 1806 1807 1808 1809 1810 1811 1812 1813 1814 1815 1816 1817 1818 1819 1820 1821 1822 1823 1824 1825 1826 1827 1828 1829 1830 1831 1832 1833 1834 1835 1836 1837 1838 1839 1840 1841 1842 1843 1844 1845 1846 1847 1848 1849 1850 1851 1852 1853 1854 1855 1856 1857 1858 1859 1860 1861 1862 1863 1864 1865 1866 1867 1868 1869 1870 1871 1872 1873 1874 1875 1876 1877 1878 1879 1880 1881 1882 1883 1884 1885 1886 1887 1888 1889 1890 1891 1892 1893 1894 1895 1896 1897 1898 1899 1900 1901 1902 1903 1904 1905 1906 1907 1908 1909 1910 1911 1912 1913 1914 1915 1916 1917 1918 1919 1920 1921 1922 1923 1924 1925 1926 1927 1928 1929 1930 1931 1932 1933 1934 1935 1936 1937 1938 1939 1940 1941 1942 1943 1944 1945 1946 1947 1948 1949 1950 1951 1952 1953 1954 1955 1956 1957 1958 1959 1960 1961 1962 1963 1964 1965 1966 1967 1968 1969 1970 1971 1972 1973 1974 1975 1976 1977 1978 1979 1980 1981 1982 1983 1984 1985 1986 1987 1988 1989 1990 1991 1992 1993 1994 1995 1996 1997 1998 1999 2000 2001 2002 2003 2004 2005 2006 2007 2008 2009 2010 2011 2012 2013 2014 2015 2016 2017 2018 2019 2020 2021 2022 2023 2024 2025 2026 2027 2028 2029 2030 2031 2032 2033 2034 2035 2036 2037 2038 2039 2040 2041 2042 2043 2044 2045 2046 2047 2048 2049 2050 2051 2052 2053 2054 2055 2056 2057 2058 2059 2060 2061 2062 2063 2064 2065 2066 2067 2068 2069 2070 2071 2072 2073 2074 2075 2076 2077 2078 2079 2080 2081 2082 2083 2084 2085 2086 2087 2088 2089 2090 2091 2092 2093 2094 2095 2096 2097 2098 2099 2100 2101 2102 2103 2104 2105 2106 2107 2108 2109 2110 2111 2112 2113 2114 2115 2116 2117 2118 2119 2120 2121 2122 2123 2124 2125 2126 2127 2128 2129 2130 2131 2132 2133 2134 2135 2136 2137 2138 2139 2140 2141 2142 2143 2144 2145 2146 2147 2148 2149 2150 2151 2152 2153 2154 2155 2156 2157 2158 2159 2160 2161 2162 2163 2164 2165 2166 2167 2168 2169 2170 2171 2172 2173 2174 2175 2176 2177 2178 2179 2180 2181 2182 2183 2184 2185 2186 2187 2188 2189 2190 2191 2192 2193 2194 2195 2196 2197 2198 2199 2200 2201 2202 2203 2204 2205 2206 2207 2208 2209 2210 2211 2212 2213 2214 2215 2216 2217 2218 2219 2220 2221 2222 2223 2224 2225 2226 2227 2228 2229 2230 2231 2232 2233 2234 2235 2236 2237 2238 2239 2240 2241 2242 2243 2244 2245 2246 2247 2248 2249 2250 2251 2252 2253 2254 2255 2256 2257 2258 2259 2260 2261 2262 2263 2264 2265 2266 2267 2268 2269 2270 2271 2272 2273 2274 2275 2276 2277 2278 2279 2280 2281 2282 2283 2284 2285 2286 2287 2288 2289 2290 2291 2292 2293 2294 2295 2296 2297 2298 2299 2300 2301 2302 2303 2304 2305 2306 2307 2308 2309 2310 2311 2312 2313 2314 2315 2316 2317 2318 2319 2320 2321 2322 2323 2324 2325 2326 2327 2328 2329 2330 2331 2332 2333 2334 2335 2336 2337 2338 2339 2340 2341 2342 2343 2344 2345 2346 2347 2348 2349 2350 2351 2352 2353 2354 2355 2356 2357 2358 2359 2360 2361 2362 2363 2364 2365 2366 2367 2368 2369 2370 2371 2372 2373 2374 2375 2376 2377 2378 2379 2380 2381 2382 2383 2384 2385 2386 2387 2388 2389 2390 2391 2392 2393 2394 2395 2396 2397 2398 2399 2400 2401 2402 2403 2404 2405 2406 2407 2408 2409 2410 2411 2412 2413 2414 2415 2416 2417 2418 2419 2420 2421 2422 2423 2424 2425 2426 2427 2428 2429 2430 2431 2432 2433 2434 2435 2436 2437 2438 2439 2440 2441 2442 2443 2444 2445 2446 2447 2448 2449 2450 2451 2452 2453 2454 2455 2456 2457 2458 2459 2460 2461 2462 2463 2464 2465 2466 2467 2468 2469 2470 2471 2472 2473 2474 2475 2476 2477 2478 2479 2480 2481 2482 2483 2484 2485 2486 2487 2488 2489 2490 2491 2492 2493 2494 2495 2496 2497 2498 2499 2500 2501 2502 2503 2504 2505 2506 2507 2508 2509 2510 2511 2512 2513 2514 2515 2516 2517 2518 2519 2520 2521 2522 2523 2524 2525 2526 2527 2528 2529 2530 2531 2532 2533 2534 2535 2536 2537 2538 2539 2540 2541 2542 2543 2544 2545 2546 2547 2548 2549 2550 2551 2552 2553 2554 2555 2556 2557 2558 2559 2560 2561 2562 2563 2564 2565 2566 2567 2568 2569 2570 2571 2572 2573 2574 2575 2576 2577 2578 2579 2580 2581 2582 2583 2584 2585 2586 2587 2588 2589 2590 2591 2592 2593 2594 2595 2596 2597 2598 2599 2600 2601 2602 2603 2604 2605 2606 2607 2608 2609 2610 2611 2612 2613 2614 2615 2616 2617 2618 2619 2620 2621 2622 2623 2624 2625 2626 2627 2628 2629 2630 2631 2632 2633 2634 2635 2636 2637 2638 2639 2640 2641 2642 2643 2644 2645 2646 2647 2648 2649 2650 2651 2652 2653 2654 2655 2656 2657 2658 2659 2660 2661 2662 2663 2664 2665 2666 2667 2668 2669 2670 2671 2672 2673 2674 2675 2676 2677 2678 2679 2680 2681 2682 2683 2684 2685 2686 2687 2688 2689 2690 2691 2692 2693 2694 2695 2696 2697 2698 2699 2700 2701 2702 2703 2704 2705 2706 2707 2708 2709 2710 2711 2712 2713 2714 2715 2716 2717 2718 2719 2720 2721 2